



CrossMark
click for updates

Report

Cite this article: Fagerlund R, Behar M, Fortmann KT, Lin YE, Vargas JD, Hoffmann A. 2015 Anatomy of a negative feedback loop: the case of $\text{I}\kappa\text{B}\alpha$. *J. R. Soc. Interface* **12**: 20150262.
<http://dx.doi.org/10.1098/rsif.2015.0262>

Received: 25 March 2015

Accepted: 3 August 2015

Subject Areas:

systems biology

Keywords:

negative feedback, gene regulation, $\text{NF}\kappa\text{B}$, $\text{I}\kappa\text{B}\alpha$, oscillation

Author for correspondence:

Alexander Hoffmann

e-mail: ahoffmann@ucla.edu

[†]Present address: Department of Biomedical Engineering, The University of Texas at Austin, Austin, TX 78712, USA.

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsif.2015.0262> or via <http://rsif.royalsocietypublishing.org>.

Anatomy of a negative feedback loop: the case of $\text{I}\kappa\text{B}\alpha$

Riku Fagerlund¹, Marcelo Behar^{1,†}, Karen T. Fortmann¹, Y. Eason Lin^{1,2,3}, Jesse D. Vargas^{1,2,3} and Alexander Hoffmann^{1,2,3}

¹Department of Chemistry and Biochemistry, University of California, San Diego, La Jolla, CA 92093, USA

²Department of Microbiology, Immunology, and Molecular Genetics, and ³Institute for Quantitative and Computational Biosciences, University of California, Los Angeles, CA 90095, USA

The magnitude, duration and oscillation of cellular signalling pathway responses are often limited by negative feedback loops, defined as an ‘activator-induced inhibitor’ regulatory motif. Within the $\text{NF}\kappa\text{B}$ signalling pathway, a key negative feedback regulator is $\text{I}\kappa\text{B}\alpha$. We show here that, contrary to current understanding, $\text{NF}\kappa\text{B}$ -inducible expression is not sufficient for providing effective negative feedback. We then employ computational simulations of $\text{NF}\kappa\text{B}$ signalling to identify $\text{I}\kappa\text{B}\alpha$ molecular properties that are critical for proper negative feedback control and test the resulting predictions in biochemical and single-cell live-imaging studies. We identified nuclear import and nuclear export of $\text{I}\kappa\text{B}\alpha$ and the $\text{I}\kappa\text{B}\alpha$ – $\text{NF}\kappa\text{B}$ complex, as well as the free $\text{I}\kappa\text{B}\alpha$ half-life, as key determinants of post-induction repression of $\text{NF}\kappa\text{B}$ and the potential for subsequent reactivation. Our work emphasizes that negative feedback is an emergent systems property determined by multiple molecular and biophysical properties in addition to the required ‘activator-induced inhibitor’ relationship.

1. Introduction

Negative feedback control is a ubiquitous regulatory motif in many biological systems, critical to the maintenance of proper homeostasis, dynamic control in response to perturbations, or oscillatory patterns [1]. The defining feature of a negative feedback motif is an activator–inhibitor pair in which the activator induces expression or activity of the inhibitor. Indeed, many studies focus on the molecular mechanism(s) that provide(s) inducibility, often characterized by the fold change and any intrinsic delay. However, actual molecular circuits within cells are incompletely described by the activator-inducible inhibitor paradigm, as they may need to contend with physical realities within the cell such as the biochemistry of molecular interactions, sub-cellular compartmentalization or protein half-life. Thus, proper functioning of a negative feedback circuit may depend on biochemical properties other than the activator-responsive control of the inhibitor.

$\text{I}\kappa\text{B}\alpha$ is a prominent negative feedback regulator in the $\text{NF}\kappa\text{B}$ signalling system [2,3]. $\text{I}\kappa\text{B}\alpha$ directly controls the dynamics of the transcription factor $\text{NF}\kappa\text{B}$, a central regulator of inflammatory and immune response gene expression [4,5]. Through its reversible sequestration of $\text{NF}\kappa\text{B}$ in the cytoplasm, $\text{I}\kappa\text{B}\alpha$ not only controls the duration of $\text{NF}\kappa\text{B}$ activity [4,6] but also enables reactivation that can result in oscillatory dynamics observed both in population studies [4] and in single cells [7]. Mathematical models were shown to recapitulate these dynamic features [8,9], and reduced models have identified $\text{NF}\kappa\text{B}$ -responsive expression of $\text{I}\kappa\text{B}\alpha$ as a key determinant of oscillatory dynamics [10–12]. The dynamics of $\text{NF}\kappa\text{B}$ signalling are stimulus-specific, and a critical determinant of inflammatory and immune gene expression programmes [13,14], prompting pioneering work to focus drug-targeting strategies on dynamical features to achieve superior specificity [15].

Here, we examine the molecular properties that confer $\text{I}\kappa\text{B}\alpha$ ’s ability to control $\text{NF}\kappa\text{B}$ dynamics. We find that while inducible expression of $\text{I}\kappa\text{B}\alpha$ is required for proper $\text{NF}\kappa\text{B}$ dynamics [16], inducible expression is not sufficient as inducible expression of another $\text{I}\kappa\text{B}$ family member, $\text{I}\kappa\text{B}\beta$, is unable to




Figure 1. (Caption overlaid.)

support normal dynamical control of NFκB. This finding prompts us to characterize other IκBα properties that are required for proper negative feedback control of NFκB. Our

study delineates how several molecular properties combine to produce the emergent systems property of dynamic negative feedback control of NFκB.

Figure 1. (*Overleaf*) NF κ B-dependent transcriptional control is not sufficient for I κ B negative functions. (a) Schematic diagram of pBabe and 5 κ B retroviral expression constructs consisting of a tandem repeat of 5 κ B sites driving the expression of I κ B α . (b) Electrophoretic mobility shift assay (EMSA) and immunoblot analysis of wt, I κ B α ^{-/-}, I κ B α ^{-/-} + pBabe_I κ B α , and I κ B α ^{-/-} + 5 κ B_I κ B α murine embryo fibroblast (MEF) cell lines. EMSA indicates NF κ B activity over a 120 min time course after stimulation with 1 ng ml⁻¹ of TNF; NFY binding was used as an EMSA control. Western blot shows protein abundances for I κ B α , I κ B β and I κ B ϵ with α -tubulin as a loading control. (c) I κ B α ^{-/-} MEFs reconstituted with NF κ B-inducible I κ B α or I κ B β were treated with 1 ng ml⁻¹ of TNF and nuclear extracts analysed by EMSA for NF κ B binding activity; NFY binding was used as an EMSA control. Immunoblots of corresponding cytoplasmic fractions were probed with antibodies specific for I κ B α , I κ B β and I κ B ϵ ; α -tubulin was used as a loading control. Densitometric quantification of NF κ B is presented as a bar graph below. (d) EMSA and immunoblot analysis of wild-type MEFs and MEFs that have the endogenous coding region for I κ B α replaced by I κ B β knock-in (AKBI). Cells were treated with 1 ng ml⁻¹ of TNF and nuclear extracts analysed by EMSA for NF κ B-binding activity; NFY binding was used as an EMSA control. Immunoblots of corresponding cytoplasmic fractions were probed with antibodies specific for I κ B α , I κ B β and I κ B ϵ ; α -tubulin was used as a loading control. Densitometric quantification of NF κ B is presented as a bar graph below. (e,f) NF κ B nuclear localization at the single-cell level. I κ B α ^{-/-} β ^{-/-} ϵ ^{-/-} RelA^{-/-} MEFs reconstituted with AcGFP1-RelA and NF κ B-inducible I κ Bs. (e) I κ B α cells were treated with 10 ng of TNF and fluorescent images were captured every 5 min and cellular localization of AcGFP1-RelA was measured and plotted as a normalized nuclear to cytoplasmic ratio individually (bottom left colour traces) and as a combined average (bottom right black trace). (f) I κ B β cells were treated with 10 ng of TNF and fluorescent images were captured every 5 min and cellular localization of AcGFP1-RelA was measured and plotted as a normalized nuclear to cytoplasmic ratio individually (bottom left colour traces) and as a combined average (bottom right black trace).

2. Results

2.1. NF κ B-responsive transcriptional control is necessary but not sufficient for I κ B α negative feedback

Studies of NF κ B dynamic control by I κ B family members have identified I κ B α as the key negative feedback regulator due to its highly inducible NF κ B-responsive promoter [2–4]. To characterize the role of NF κ B-inducible expression, we complemented I κ B α -deficient murine embryo fibroblasts (MEFs) with retroviral plasmids that express I κ B α from either a constitutive (pBabe) or an NF κ B-inducible (5 κ B) promoter (figure 1a). Unlike pBabe-reconstituted cells, 5 κ B_I κ B α reconstituted cells showed dynamic resynthesis profiles similar to endogenous I κ B α in wild-type cells following stimulation with tumour necrosis factor (TNF) (figure 1b). Importantly, when we examined the control of NF κ B activity by electrophoretic mobility shift assay (EMSA), we found that 5 κ B cells showed post-induction repression and the transient trough of NF κ B activity characteristic of wild-type cells, correcting the misregulation in I κ B α -deficient cells, whereas cells constitutively expressing I κ B α were unable to capture this response (figure 1b).

To test whether NF κ B-inducible control was not only required but also sufficient for NF κ B dynamic control, we complemented I κ B α -deficient cells with a 5 κ B retrovirus expressing I κ B β , a highly homologous I κ B family member capable of inhibiting NF κ B but not normally providing negative feedback. Interestingly, these cells did not show proper dynamic control of NF κ B even though I κ B β expression was under NF κ B control similar to I κ B α (figure 1c; electronic supplementary material, figure S1A). However, when another known I κ B negative feedback regulator, I κ B ϵ [17], was linked to this promoter, NF κ B activity did show post-induction repression (electronic supplementary material, figure S1B). These results indicate that, despite the high degree of sequence homology, I κ B α and I κ B β have distinct molecular properties that, along with differential gene expression control, render I κ B α an effective negative feedback regulator but not I κ B β . In order to confirm the validity of this conclusion, we obtained fibroblasts from a genetic knock-in mouse in which the I κ B β coding region was engineered to replace the I κ B α open reading frame such that I κ B β expression was under the control of the endogenous I κ B α promoter [18]. Remarkably, these so-called AKBI cells also failed to show proper NF κ B post-induction

attenuation despite highly inducible I κ B β expression (figure 1d; electronic supplementary material, figure S1c).

In order to examine translocation dynamics in single cells, and without the confounding contributions of other I κ B family members, we generated I κ B α ^{-/-} β ^{-/-} ϵ ^{-/-} RelA^{-/-} T3 cells that lack all three classical NF κ B inhibitors and RelA, and reconstituted them with a constitutively expressed fluorescent GFP-RelA and NF κ B-responsively expressed I κ B α or I κ B β . Whereas reconstitution with I κ B α resulted in transient NF κ B activation in response to TNF treatment, defined by a trough at about 60 min, followed by a second phase in some cells (figure 1e), reconstitution with NF κ B-inducible I κ B β resulted in sustained NF κ B activation showing only slow and incomplete post-induction repression (figure 1f). Of note, in both conditions, the mean RelA nuclear localization profile of the collection of individual cells (figure 1e,f, black trace) closely resembled the population level in biochemical studies (figure 1c,d). These data clearly indicate that when expression of I κ B α or I κ B β is driven by the same NF κ B-responsive promoter, resulting in ostensibly similar expression profiles, only I κ B α can provide effective dynamic negative feedback control on NF κ B. Thus, inducible inhibitor expression in and of itself is not sufficient for proper negative feedback control of NF κ B.

2.2. Mathematical modelling identifies multiple molecular properties of I κ B α contributing to the negative feedback control of NF κ B

I κ B α has several characteristics—other than NF κ B-dependent synthesis—that in principle may contribute to its negative feedback function, e.g. its nuclear import and export properties, as well as constitutive and signal-induced degradation of free and NF κ B-bound I κ B α (figure 2a). Here, we use a previously established *in silico* model of NF κ B regulation to investigate the contributions of each of these processes to the control of dynamic NF κ B signals. When normalized for maximum activity, we confirmed that partial inhibition of the NF κ B-dependent synthesis of I κ B α potently impaired post-induction attenuation, with 10% inhibition resulting in a 30% increase in the signalling level at 70 min (figure 2b, row 1, and 2c). However, we also found that partial inhibition of I κ B α nuclear import had a similar effect with a 10% inhibition causing an 11% increase in signalling at 70 min (figure 2b, row 3). Weak inhibition of the degradation of free I κ B α had little effect on post-attenuation induction, whereas stronger inhibition shifted




Figure 2. Modelling IκBα properties contributing to negative feedback control of NFκB. (a) Schematic illustrates negative feedback control of NFκB by IκBs. Numbers indicate potential reactions that may contribute to dynamic regulation of NFκB. (b) Normalized nuclear NFκB concentration is shown for unperturbed models (black) and models in which the indicated reactions are partially inhibited (blue-red lines reflect various degrees of inhibition). Reaction numbers as in (a). (c) Sensitivity analysis of three temporal phases of the NFκB with respect to changes in IκBα regulation. (i)–(iii) Sensitivity ratio $(\text{nucNF}\kappa\text{B}^{\text{perturbed}} - \text{nucNF}\kappa\text{B}^{\text{unperturbed}}) / \text{nucNF}\kappa\text{B}^{\text{unperturbed}}$ in per cent units for each reaction (numbers as in (a)). (iv) Global sensitivity (RMSD) integrated over 120 min. Results are shown for three levels of inhibition (10%, twofold and 10-fold).

the peak of NFκB to later times resulting in a modest increase in late activity (90% inhibition caused 14% increase in signalling at 120 min, figure 2b, row 2). Partial inhibition of nuclear export or of signal-induced degradation of NFκB-bound IκBα also reduced the post-attenuation reactivation, with a 10% inhibition causing 1% and 9.5% decreased signalling at 120 min, respectively (figure 2b, rows 4 and 5).

To compare the contribution of these processes, we determined the sensitivity of NFκB activity at three specific times representing early, post-induction attenuation and late parts of the signal to various perturbations (figure 2c). We also quantified the global sensitivity to each perturbation as the root mean square deviation (RMSD) of the perturbed and unperturbed signals over 120 min, sampled at 1 min intervals

(figure 2c(iv)). This analysis posits that IκBα-mediated post-induction attenuation of NFκB activity (figure 2c(ii)) is a function not only of the NFκB-dependent synthesis rate but also of IκBα's nuclear import, as well as its constitutive degradation. It also predicts that nuclear export and IKK-dependent degradation of NFκB-bound IκBα are important for late post-attenuation signalling (figure 2c(iii)).

2.3. Experimental testing of model predictions: multiple IκBα properties contribute distinct characteristics to NFκB dynamic control

To test the computational predictions, we pursued a genetic perturbation approach. Previous work showed nuclear

import of $\text{I}\kappa\text{B}\alpha$ to be mediated by an unconventional NLS sequence [19,20]. Using this information, we reconstituted $\text{I}\kappa\text{B}\alpha$ -deficient cells with an $\text{I}\kappa\text{B}\alpha$ NLS mutant ($\text{I}\kappa\text{B}\alpha\text{NLSm}$: L110A,L115A,L117A,L120A). In DNA binding studies of $\text{I}\kappa\text{B}\alpha\text{NLSm}$ cells, TNF induced NF κ B activation comparable to that of wild-type $\text{I}\kappa\text{B}\alpha$ cells (figure 3a), and although the resynthesis of $\text{I}\kappa\text{B}\alpha\text{NLSm}$ protein was effectively induced by NF κ B, the $\text{I}\kappa\text{B}\alpha\text{NLSm}$ cells were defective for the rapid post-induction repression of NF κ B. At 70 min, the signal in $\text{I}\kappa\text{B}\alpha\text{NLSm}$ cells was 2.9-fold higher than in wild-type $\text{I}\kappa\text{B}\alpha$ cells (considering the different basal levels). This is consistent with the threefold increase predicted by the model when the corresponding parameter is reduced to 35% of its wild-type value (figure 3a; electronic supplementary material, figure S2a). Consistent with population-level biochemical studies, $\text{I}\kappa\text{B}\alpha^{-/-}\beta^{-/-}\epsilon^{-/-}\text{RelA}^{-/-}$ cells expressing GFP-RelA showed that $\text{I}\kappa\text{B}\alpha\text{NLSm}$ was defective in post-induction repression and cytoplasmic relocalization of NF κ B in single cells (figure 3b). Despite the substantial heterogeneity in RelA cytoplasmic re-localization, the population mean closely resembles the population-level results obtained by EMSA. These data clearly show that the nuclear localization of $\text{I}\kappa\text{B}\alpha$ is indispensable for proper termination of NF κ B activity.

$\text{I}\kappa\text{B}\alpha$ also relies on a nuclear export sequence (NES) for the efficient nuclear export of NF κ B [21,22]. Mathematical modelling suggested strong inhibition of nuclear export would result in reduced late NF κ B activity. To assess the role of $\text{I}\kappa\text{B}\alpha$ nuclear export on the sub-cellular localization control of NF κ B, we generated an NES mutant ($\text{I}\kappa\text{B}\alpha\text{NESm}$; L45A,L49A,I52A). Reconstituted cells expressing $\text{I}\kappa\text{B}\alpha\text{NESm}$ were unable to produce the post-repression reactivation of NF κ B characteristic of wild-type protein-controlled NF κ B signalling (figure 3c; electronic supplementary material, figure S2b). The activity is qualitatively similar to the model prediction for a fivefold attenuation in the corresponding parameter. These results indicate that the nuclear export function of $\text{I}\kappa\text{B}\alpha$ is crucial for the post-repression activation of NF κ B signalling. Interestingly, the single-cell studies with the NES mutant revealed seemingly contradictory results (figure 3d), as most cells displayed sustained RelA nuclear localization. This apparent discrepancy is resolved by recognizing that: (1) the mutant localizes to the nucleus causing inhibition of NF κ B activity but does not allow for NF κ B export and reactivation, and (2) the biochemical assay detects DNA binding activity of free NF κ B, whereas the single-cell imaging is a readout of NF κ B localization only.

$\text{I}\kappa\text{B}\alpha$ is known to have a very short half-life that is extended approximately twofold by an $\text{I}\kappa\text{B}\alpha 5\text{M}$ mutant (S283A, S288,T291A,S293A,T296A) [23,24]. When expressed in $\text{I}\kappa\text{B}\alpha^{-/-}$ cells from an NF κ B-responsive promoter, $\text{I}\kappa\text{B}\alpha 5\text{M}$ achieved effective post-induction repression of NF κ B activity (figure 3e; electronic supplementary material, figure S2c). However, the re-activation of NF κ B was undetectable, consistent with computational predictions that indicated a signal close to basal level at 120 min when the corresponding parameter was reduced to 35% of its wild-type value. Similarly, in single live-cell studies, $\text{I}\kappa\text{B}\alpha 5\text{M}$ mediated efficient relocalization of RelA to the cytoplasm with a complete absence of late-phase activity (figure 3f).

3. Discussion

Given the well-documented role of NF κ B activity in vital cellular processes, a number of mechanisms have evolved

to ensure precise regulation of its activity. The $\text{I}\kappa\text{B}\alpha$ negative feedback loop is a prominent NF κ B regulatory mechanism, allowing for both post-induction repression and repeated or oscillatory bursts of activity, and is critical for providing complex dynamic control which is thought to mediate specificity in NF κ B's pleiotropic physiological functions. Prior studies have established that the NF κ B-responsive $\text{I}\kappa\text{B}\alpha$ promoter is critical for this negative feedback control [16], but it has remained unclear whether specific characteristics of the $\text{I}\kappa\text{B}\alpha$ protein may be important as well. Indeed, biochemical studies presented here using MEFs derived from mice in which $\text{I}\kappa\text{B}\beta$ was engineered into the $\text{I}\kappa\text{B}\alpha$ locus to (AKBI MEFs) clearly demonstrate that, even when $\text{I}\kappa\text{B}\beta$ is under NF κ B-transcriptional induction, it is unable to provide proper negative feedback. These data motivated our characterization of $\text{I}\kappa\text{B}\alpha$ protein properties that contribute to proper negative feedback function. Our strategy was to complement $\text{I}\kappa\text{B}\alpha^{-/-}$ cells with retroviral transgenes providing for κ B-responsive expression of engineered $\text{I}\kappa\text{B}\alpha$ variants defective in specific molecular characteristics.

In addition to traditional biochemical approaches to study NF κ B response and regulation, we examined NF κ B response dynamics in single cells. Recent studies have characterized NF κ B dynamics in single cells, but, to date, no studies have employed gene knock-out cells to probe underlying molecular mechanisms. Thus, regulatory control mechanisms identified at the biochemical/population level have yet to be reconciled with single-cell microscopy tracking studies that boast high temporal resolution and individual cellular histories. In this work, we employed a cell line lacking RelA and all canonical $\text{I}\kappa\text{B}$ proteins ($\text{I}\kappa\text{B}\alpha^{-/-}\text{I}\kappa\text{B}\beta^{-/-}\text{I}\kappa\text{B}\epsilon^{-/-}\text{RelA}^{-/-}$ cells), which we then reconstituted with NF κ B-inducible $\text{I}\kappa\text{B}$ variants and fluorescent RelA reporter in order to examine the contributions of specific $\text{I}\kappa\text{B}$ protein characteristics.

Although the $\text{I}\kappa\text{B}$ proteins were first identified as cytoplasmic inhibitors, it has become clear that they play a major role in regulating nuclear NF κ B. $\text{I}\kappa\text{B}\alpha$ has been shown to be efficiently transported into the nucleus where it binds active NF κ B dimers on the promoters of NF κ B-activated genes and facilitates the dissociation of the transcription factor from DNA. Comparing mutants with wild-type $\text{I}\kappa\text{B}$ proteins, we were able to show the contributions of $\text{I}\kappa\text{B}$ inducible synthesis, nucleo-cytoplasmic transport and degradation control to the various aspects of the prototypical NF κ B response; namely, duration and amplitude of initial NF κ B activation, post-activation repression and post-repression re-activation of NF κ B signalling. Our biochemical assays, together with single-cell studies, demonstrated that $\text{I}\kappa\text{B}\alpha$ nuclear localization is indispensable for the rapid termination of NF κ B activity. Specifically, we showed that an NES-deficient form of $\text{I}\kappa\text{B}\alpha$ supported efficient induction and post-induction repression of NF κ B DNA binding activity, but not the characteristic re-activation and second phase NF κ B activity. The deficiency in NES function prevents the protein from efficiently returning NF κ B to the cytoplasm for the next round of activation, maintaining an inactive $\text{I}\kappa\text{B}\alpha\text{NESm}$ -bound pool of NF κ B in the nucleus. Finally, by employing an $\text{I}\kappa\text{B}\alpha$ harbouring five mutations that confer stability, increasing the half-life of the normally rapidly turned over uncomplexed/free protein, we were able to show the importance of such rapid turnover in generating characteristic NF κ B temporal profiles. In cells




Figure 3. Multiple $I\kappa B\alpha$ properties contribute distinct characteristics to NFκB control. Nuclear localization of $I\kappa B\alpha$ is required for the termination of NFκB activity (*a,b*). Nuclear export function of $I\kappa B\alpha$ is required for post-repression activation of NFκB activity (*c,d*). $I\kappa B\alpha$ protein half-life control is critical for sustained NFκB dynamics (*e,f*). $I\kappa B\alpha^{-/-}$ MEFs were reconstituted with NFκB-inducible wild-type or NLS mutant (*a,b*), NES mutant (*c,d*) or the 5M mutant (*e,f*) form of $I\kappa B\alpha$. The cells were treated with 1 ng ml^{-1} of TNF and nuclear extracts were analysed by EMSA for NFκB activity and corresponding cytoplasmic extracts subjected to western blotting with indicated antibodies (*a,c,e*). Bar graphs show quantification of EMSA; curves are modelling the result of NFκB activity in single cells upon stimulation with 10 ng ml^{-1} of TNF. Real-time fluorescent images of $I\kappa B\alpha^{-/-} \beta^{-/-} \epsilon^{-/-} RelA^{-/-}$ MEFs reconstituted with AcGFP1-RelA and NFκB-inducible $I\kappa B\alpha$ NLS mutant (*b*), NES mutant (*d*) or the 5M mutant (*f*) $I\kappa B\alpha$ (showing cellular localization of RelA at indicated time points). Below the fluorescent images, single-cell traces show the ratio of nuclear to cytoplasmic localization of AcGFP1-RelA in fluorescent images (left) as well as the average curve and standard deviation of the single-cell traces (right).

expressing this $I\kappa B\alpha 5M$ form, we found a complete absence of second-phase activation, indicating that a low level of free $I\kappa B\alpha$ protein (ensured by a short half-life) is required for this aspect of the response.

Our results demonstrate not only that $I\kappa B\alpha$ feedback is dependent on $NF\kappa B$ -inducible synthesis but also that several other processes dependent on the molecular characteristics of the protein itself, for example import, export and half-life control, must be tuned in a coordinated manner to generate the hallmark features of $NF\kappa B$ signaling, namely post-induction repression and reactivation. By contrast, the $I\kappa B\beta$ protein does not support proper negative feedback control even when expressed from $I\kappa B\alpha$'s promoter; we speculate that substantially reduced nucleocytoplasmic transport [25,26] may be a key underlying reason; a second characteristic that may play a role is $I\kappa B\alpha$'s but not $I\kappa B\beta$'s ability to strip $NF\kappa B$ off the DNA [27]. Indeed, these properties are not required for the $NF\kappa B$ dimer stabilization/chaperone function recently ascribed to $I\kappa B\beta$ [28]. Our results illustrate the more general point: that negative feedback regulation in cells is a complex process that depends on multiple molecular properties beyond activator-induced expression [29,30]. These findings may well extend to other transcriptional networks as nuclear transport is a defining feature of many gene regulatory networks. Understanding the specific contributions of each process as well as their characteristic time scales is an important step for identifying effective druggable targets that may allow for correction of dynamic misregulation in cells associated with pathology [15].

4. Material and methods

4.1. Computational modelling

The response of the $NF\kappa B$ regulatory module was simulated using the computational ODE-based model described in [31]. In order to focus on regulatory mechanisms involving $I\kappa B\alpha$, the other $I\kappa B$ family members were removed. Following equilibration, TNF responses were simulated as in [31]. Time-course curves in figure 2 were generated by applying multipliers to the kinetic parameters corresponding to the reactions in figure 2a. The multiplier values were: $2^{-3}, -2.5, -2, -1.5, -1, -0.5$ (reactions 1, 2, 3 and 5) and $2^{-6}, -5, -4, -3, -2, -1$ (reaction 4), reflecting different sensitivities for reaction 4. $NF\kappa B$ time courses are normalized to their peak value. Sensitivity ratios $s_r(t)$ at a particular time t_i are defined as: $(nucNF\kappa B^{perturbed} - nucNF\kappa B^{unperturbed})/nucNF\kappa B^{unperturbed}$, where $nucNF\kappa B^{perturbed/unperturbed}$ are the normalized nuclear concentrations of $NF\kappa B$ at time t_i obtained with a model with/without a multiplicative factor (0.9, 0.5, 0.1) for the indicated kinetic rate parameter (values shown in per cent units). The global average sensitivity in figure 2c was calculated as the RMS of the $s_r(t)$ sampled at 1 min intervals between 1 and 120 min post-stimulation.

4.2. DNA constructs

$NF\kappa B$ -inducible $I\kappa B\alpha$ and $I\kappa B\beta$ constructs were generated in the self-inactivating (SIN) retrovirus backbone (HRSpuro) modified to express the $I\kappa B\alpha$ or $I\kappa B\beta$ transgene under the control of five tandem κB sites upstream of a minimal promoter. $I\kappa B\alpha$ mutant forms were produced using site-directed mutagenesis. For live-cell studies, AcGFP1 was fused to the N-terminus of RelA and the resulting construct was sub-cloned into the constitutively expressing retroviral plasmid pBabe-Hygro.

4.3. Cells and cell culture

Immortalized $I\kappa B\alpha^{-/-}$ MEFs were previously described [4] and $I\kappa B\alpha^{-/-}\beta^{-/-}\epsilon^{-/-}RelA^{-/-}$ MEFs were produced by interbreeding of the four individual mouse knock-out strains and harvesting E13.5 embryos, subjecting primary MEFs to the 3T3 protocol of repeated passage until a stably proliferating cell culture emerged. AKBI MEFs were a generous gift from BingBing Jiang (Boston University). MEFs were cultured in Dulbecco's modified Eagle's medium supplemented with 100 U penicillin/streptomycin (10378016; Life Technologies), 0.3 mg ml⁻¹ glutamine and 10% fetal calf serum (complete medium). Plat-E cells [32] were maintained in complete medium containing blasticidin (10 µg ml⁻¹) and puromycin (1 µg ml⁻¹).

4.4. Retrovirus-mediated gene transduction

$NF\kappa B$ -inducible $I\kappa B$ and AcGFP1-RelA constructs were transfected into Plat-E packaging cells pre-conditioned in antibiotic-free complete medium using poly(ethylenimine). Supernatant was collected 48 h post-transfection, filtered and used to infect target cells with 4 µg ml⁻¹ polybrene to enhance infection efficiency (Sigma). Infected cells were selected with puromycin hydrochloride (Sigma) for $I\kappa Bs$ and/or with hygromycin B (InvivoGen) for the AcGFP1-RelA. Murine TNF (Roche) was used at 1 or 10 ng ml⁻¹.

4.5. Biochemical analyses

Whole-cell extracts were prepared in radioimmunoprecipitation assay buffer with protease inhibitors and normalized for total protein before immunoblot analyses. Cytoplasmic and nuclear extracts for immunoblot analyses and EMSA, respectively, were prepared as previously described [4,24]. $I\kappa B\alpha$ was probed with sc-371, $I\kappa B\beta$ with sc-945 and α -tubulin with sc-5286. All antibodies were from Santa Cruz Biotechnology.

4.6. Microscopy

Cells were plated onto 35 mm glass bottom dishes (MatTek) or iBidi eight-well chambers (iBidi) 24 h prior to stimulation and immediate imaging. Images were acquired on an Axio Observer Z1 inverted microscope (Carl Zeiss Microscopy GmbH, Germany) with a 40 \times , 1.3 NA oil-immersion, or 20 \times , 0.8 NA air-immersion objective to a Coolsnap HQ2 CCD camera (Photometrics, Canada) using ZEN imaging software (Carl Zeiss Microscopy GmbH, Germany). Environmental conditions were maintained in a humidified chamber at 37°C, 5% CO₂ (Pecon, Germany). Quantitative image processing was performed using the FIJI distribution of IMAGE J (NIH). All cells of each frame in the microscope imaging experiments were measured for total fluorescence intensity. Time-course data were normalized by the minimum and maximum values to account for the varying overall intensities of different cells. The single-cell traces were averaged and error bars in the mean curves are the standard deviation from the mean.

Authors' contributions. R.F. performed the experimental work, assisted by K.T.F., Y.E.L. and J.D.V. M.B. performed the computational modelling. R.F., J.D.V. and A.H. wrote the manuscript.

Competing interests. We have no competing interests.

Funding. The work was supported by grants to A.H. from the NIH: R01 GM071573 and P01 GM071862. R.F. was a Sigrid Juselius Foundation and Saatioiden postdoctoral fellow and M.B. was a Cancer Research Institute postdoctoral fellow.

Acknowledgements. We thank Bingbing Jiang (Boston University) for the generous gift of AKBI MEFs and acknowledge Santa Cruz Biotechnology for their support.

References

- Alon U. 2007 *An introduction to systems biology: design principles of biological circuits*. Boca Raton, FL: Chapman and Hall/CRC.
- Scott ML, Fujita T, Liou HC, Nolan GP, Baltimore D. 1993 The p65 subunit of NF-kappa B regulates I kappa B by two distinct mechanisms. *Genes Dev.* **7**, 1266–1276. (doi:10.1101/gad.7.7a.1266)
- Chiao PJ, Miyamoto S, Verma IM. 1994 Autoregulation of I kappa B alpha activity. *Proc. Natl Acad. Sci. USA* **91**, 28–32. (doi:10.1073/pnas.91.1.28)
- Hoffmann A, Levchenko A, Scott ML, Baltimore D. 2002 The IkappaB-NF-kappaB signaling module: temporal control and selective gene activation. *Science* **298**, 1241–1245. (doi:10.1126/science.1071914)
- Hoffmann A, Baltimore D. 2006 Circuitry of nuclear factor kappaB signaling. *Immunol. Rev.* **210**, 171–186. (doi:10.1111/j.0105-2896.2006.00375.x)
- Shih VF, Kearns JD, Basak S, Savinova OV, Ghosh G, Hoffmann A. 2009 Kinetic control of negative feedback regulators of NF-kappaB/RelA determines their pathogen- and cytokine-receptor signaling specificity. *Proc. Natl Acad. Sci. USA* **106**, 9619–9624. (doi:10.1073/pnas.0812367106)
- Nelson DE *et al.* 2004 Oscillations in NF-kappaB signaling control the dynamics of gene expression. *Science* **306**, 704–708. (doi:10.1126/science.1099962)
- O'Dea E, Hoffmann A. 2010 The regulatory logic of the NF-kappaB signaling system. *Cold Spring Harb. Perspect. Biol.* **2**, a000216. (doi:10.1101/cshperspect.a000216)
- Basak S, Behar M, Hoffmann A. 2012 Lessons from mathematically modeling the NF-kappaB pathway. *Immunol. Rev.* **246**, 221–238. (doi:10.1111/j.1600-065X.2011.01092.x)
- Krishna S, Jensen MH, Sneppen K. 2006 Minimal model of spiky oscillations in NF-kappaB signaling. *Proc. Natl Acad. Sci. USA* **103**, 10 840–10 845.
- Hayot F, Jayaprakash C. 2006 NF-kappaB oscillations and cell-to-cell variability. *J. Theor. Biol.* **240**, 583–591. (doi:10.1016/j.jtbi.2005.10.018)
- Mothes J, Busse D, Kofahl B, Wolf J. 2015 Sources of dynamic variability in NF-kappaB signal transduction: a mechanistic model. *BioEssays* **37**, 452–462. (doi:10.1002/bies.201400113)
- Werner SL, Barken D, Hoffmann A. 2005 Stimulus specificity of gene expression programs determined by temporal control of IKK activity. *Science* **309**, 1857–1861. (doi:10.1126/science.1113319)
- Behar M, Hoffmann A. 2010 Understanding the temporal codes of intra-cellular signals. *Curr. Opin. Genet. Dev.* **20**, 684–693. (doi:10.1016/j.gde.2010.09.007)
- Behar M, Barken D, Werner SL, Hoffmann A. 2013 The dynamics of signaling as a pharmacological target. *Cell* **155**, 448–461. (doi:10.1016/j.cell.2013.09.018)
- Werner SL, Kearns JD, Zadorozhnyaya V, Lynch C, O'Dea E, Boldin MP, Ma A, Baltimore D, Hoffmann A. 2008 Encoding NF-kappaB temporal control in response to TNF: distinct roles for the negative regulators IkappaBalpha and A20. *Genes Dev.* **22**, 2093–2101. (doi:10.1101/gad.1680708)
- Kearns JD, Basak S, Werner SL, Huang CS, Hoffmann A. 2006 IkappaBepsilon provides negative feedback to control NF-kappaB oscillations, signaling dynamics, inflammatory gene expression. *J. Cell Biol.* **173**, 659–664. (doi:10.1083/jcb.200510155)
- Cheng JD, Ryseck RP, Attar RM, Dambach D, Bravo R. 1998 Functional redundancy of the nuclear factor kappa B inhibitors I kappa B alpha and I kappa B beta. *J. Exp. Med.* **188**, 1055–1062. (doi:10.1084/jem.188.6.1055)
- Sachdev S, Bagchi S, Zhang DD, Mings AC, Hannink M. 2000 Nuclear import of IkappaBalpha is accomplished by a ran-independent transport pathway. *Mol. Cell Biol.* **20**, 1571–1582. (doi:10.1128/MCB.20.5.1571-1582.2000)
- Sachdev S, Hoffmann A, Hannink M. 1998 Nuclear localization of IkappaB alpha is mediated by the second ankyrin repeat: the IkappaB alpha ankyrin repeats define a novel class of cis-acting nuclear import sequences. *Mol. Cell Biol.* **18**, 2524–2534.
- Huang TT, Kudo N, Yoshida M, Miyamoto S. 2000 A nuclear export signal in the N-terminal regulatory domain of IkappaBalpha controls cytoplasmic localization of inactive NF-kappaB/IkappaBalpha complexes. *Proc. Natl Acad. Sci. USA* **97**, 1014–1019. (doi:10.1073/pnas.97.3.1014)
- Huang TT, Miyamoto S. 2001 Postrepression activation of NF-kappaB requires the amino-terminal nuclear export signal specific to IkappaBalpha. *Mol. Cell Biol.* **21**, 4737–4747. (doi:10.1128/MCB.21.14.4737-4747.2001)
- Mathes E, O'Dea EL, Hoffmann A, Ghosh G. 2008 NF-kappaB dictates the degradation pathway of IkappaBalpha. *EMBO J.* **27**, 1357–1367. (doi:10.1038/emboj.2008.73)
- O'Dea EL, Kearns JD, Hoffmann A. 2008 UV as an amplifier rather than inducer of NF-kappaB activity. *Mol. Cell* **30**, 632–641. (doi:10.1016/j.molcel.2008.03.017)
- Chen Y, Wu J, Ghosh G. 2003 KappaB-Ras binds to the unique insert within the ankyrin repeat domain of IkappaBbeta and regulates cytoplasmic retention of IkappaBbeta x NF-kappaB complexes. *J. Biol. Chem.* **278**, 23 101–23 106. (doi:10.1074/jbc.M301021200)
- Malek S, Chen Y, Huxford T, Ghosh G. 2001 IkappaBbeta, but not IkappaBalpha, functions as a classical cytoplasmic inhibitor of NF-kappaB dimers by masking both NF-kappaB nuclear localization sequences in resting cells. *J. Biol. Chem.* **276**, 45 225–45 235. (doi:10.1074/jbc.M105865200)
- Bergqvist S, Alverdi V, Mengel B, Hoffmann A, Ghosh G, Komives EA. 2009 Kinetic enhancement of NF-kappaBxDNA dissociation by IkappaBalpha. *Proc. Natl Acad. Sci. USA* **106**, 19 328–19 333.
- Tsui R, Kearns JD, Lynch C, Vu D, Ngo KA, Basak S, Ghosh G, Hoffmann A. 2015 IkappaBbeta enhances the generation of the low-affinity NFkappaB/RelA homodimer. *Nat. Commun.* **6**, 7068. (doi:10.1038/ncomms8068)
- Nguyen LK, Kulasiri D. 2009 On the functional diversity of dynamical behaviour in genetic and metabolic feedback systems. *BMC Syst. Biol.* **3**, 51. (doi:10.1186/1752-0509-3-51)
- Nguyen LK. 2012 Regulation of oscillation dynamics in biochemical systems with dual negative feedback loops. *J. R. Soc. Interface* **9**, 1998–2010. (doi:10.1098/rsif.2012.0028)
- Mukherjee SP, Behar M, Birnbaum HA, Hoffmann A, Wright PE, Ghosh G. 2013 Analysis of the RelA:CBP/p300 interaction reveals its involvement in NF-kappaB-driven transcription. *PLoS Biol.* **11**, e1001647. (doi:10.1371/journal.pbio.1001647)
- Morita S, Kojima T, Kitamura T. 2000 Plat-E: an efficient and stable system for transient packaging of retroviruses. *Gene Ther.* **7**, 1063–1066. (doi:10.1038/sj.gt.3301206)