

Calcification rates and the effect of ocean acidification on Mediterranean cold-water corals

C. Maier^{1,2,*}, P. Watremez³, M. Taviani^{4,5}, M. G. Weinbauer^{1,2}
and J. P. Gattuso^{1,2}

¹Department of Microbiology and Biogeochemistry, CNRS-INSU, Laboratoire d'Océanographie de Villefranche, BP 28, 06234 Villefranche-sur-Mer Cedex, France

²Université Pierre et Marie Curie-Paris 6, Observatoire Océanologique de Villefranche, 06230 Villefranche-sur-Mer, France

³Agence des aires marines protégées, 16 quai de la Douane, 29229 Brest, France

⁴Ismar-CNR, via Gobetti 101, 40129 Bologna, Italy

⁵Biology Department, Woods Hole Oceanographic Institution, 266 Woods Hole Road, Woods Hole, MA 02543, USA

Global environmental changes, including ocean acidification, have been identified as a major threat to scleractinian corals. General predictions are that ocean acidification will be detrimental to reef growth and that 40 to more than 80 per cent of present-day reefs will decline during the next 50 years. Cold-water corals (CWCs) are thought to be strongly affected by changes in ocean acidification owing to their distribution in deep and/or cold waters, which naturally exhibit a CaCO_3 saturation state lower than in shallow/warm waters. Calcification was measured in three species of Mediterranean cold-water scleractinian corals (*Lophelia pertusa*, *Madrepora oculata* and *Desmophyllum dianthus*) on-board research vessels and soon after collection. Incubations were performed in ambient sea water. The species *M. oculata* was additionally incubated in sea water reduced or enriched in CO_2 . At ambient conditions, calcification rates ranged between -0.01 and $0.23\% \text{ d}^{-1}$. Calcification rates of *M. oculata* under variable partial pressure of CO_2 (pCO_2) were the same for ambient and elevated pCO_2 (404 and $867 \mu\text{atm}$) with $0.06 \pm 0.06\% \text{ d}^{-1}$, while calcification was $0.12 \pm 0.06\% \text{ d}^{-1}$ when pCO_2 was reduced to its pre-industrial level ($285 \mu\text{atm}$). This suggests that present-day CWC calcification in the Mediterranean Sea has already drastically declined (by 50%) as a consequence of anthropogenic-induced ocean acidification.

Keywords: cold-water coral; *Lophelia pertusa*; *Madrepora oculata*; *Desmophyllum dianthus*; calcification; ocean acidification

1. INTRODUCTION

Cold-water corals (CWCs) have recently come into the focus of research as biodiversity hotspots in the deep oceans [1]. Although, they have been described since the early eighteenth century, their more comprehensive study has been, and still is, restricted by the fact that they predominantly occur in deep depths or on steep cliffs, canyon walls or under overhangs. Technological developments with respect to shipboard logistics, as well as acoustic and video survey techniques, has led to the discovery of a large number of CWC sites during the last decades [2–9].

CWCs are ecosystem engineers which need to outgrow sedimentation and other destructive forces in order to maintain a three-dimensional structure that serves as the foundation for the ecosystem. Skeletal extension rates between 6 and 26 mm yr^{-1} for *Lophelia pertusa* were derived from indirect measurements of stable isotopes [10–13] and from corals growing on artificial substrate [14,15]. For corals maintained in aquaria, skeletal extension rates were

9.4 mm yr^{-1} and $15\text{--}17 \text{ mm yr}^{-1}$ for *L. pertusa* from Norwegian fjords and the Mediterranean Sea, respectively [16,17], and $3\text{--}18 \text{ mm yr}^{-1}$ for *Madrepora oculata* from the Mediterranean Sea [17]. One study reported average extension rates of 3.77 mm yr^{-1} and 2.44 mm yr^{-1} for branches of *L. pertusa* stained with Alizarin Red S and re-deployed *in situ* at coral and non-coral areas, respectively [18]. Studies of CWC growth usually refer to linear or radial extension which, without information on skeletal density, provides no information on calcification rates (mass of CaCO_3 precipitated per unit time) [19]. So far, only two studies have reported rates of calcification in CWCs. Average calcification rates measured with the buoyant weighing technique ranged between 0.02 and $0.11\% \text{ d}^{-1}$ in four Mediterranean species [20]. Maier *et al.* [21] reported average rates of $0.02\text{--}0.07\% \text{ d}^{-1}$ obtained through labelling of colonies of *L. pertusa* with the radiotracer ^{45}Ca .

It is well known that the distribution of CWCs is limited by temperature rather than depth and they are most commonly distributed at temperatures between 4°C and 12°C [1]. In the Mediterranean Sea, they occur at temperatures between 12.5°C , to almost 14°C , which might already be at the higher tolerance limit [2]. Furthermore, the uptake of anthropogenic CO_2 by the ocean generates

*Author for correspondence (maier.conny@gmail.com).

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2011.1763> or via <http://rspb.royalsocietypublishing.org>.

ocean acidification which has recently been proclaimed as one of the perhaps most harmful threats to CWCs [1]. That is because CWCs are mostly distributed at relatively high latitudes and/or deep water. Both environments will be the first to be affected by the lowering in ocean pH and a shallowing of the aragonite saturation (Ω_a) horizon [22–24]. The Mediterranean Sea with its two Western and Eastern main basins, separated by the Straits of Sicily, and some smaller regional basins, can be regarded as a miniature ocean that is expected to react faster to global change than the open ocean [25]. Over the last decades, there appears to have been a warming trend attributed to global warming in deep [26] and surface waters [27,28]. A very recent study has shown that there is a high imprint of anthropogenic CO_2 , which has already affected Mediterranean pH by a decrease of 0.05–0.14 pH units since pre-industrial times [29]. Owing to the high alkalinity in the Mediterranean, more atmospheric CO_2 is absorbed than in the open ocean [30]. It is consequently important to understand how anthropogenic partial pressure of CO_2 ($p\text{CO}_2$) has already affected and how it will affect Mediterranean Sea ecosystems and its key taxa.

Because of their restricted accessibility, it is a demanding task to study the biology and physiology of CWCs, making it important to develop reproducible methods for experimentation. One option is to maintain corals under controlled laboratory conditions, and attempt to mimic their natural environment. However, the maintenance of CWCs in the laboratory is very time consuming and prone to technical problems. We therefore conducted on-board experiments with freshly collected CWCs and ambient sea water. The aim of the study was to compare rates of calcification in three Mediterranean species at three study sites and to test the effects of ocean acidification on *M. oculata* calcification rates by subjecting colonies to three $p\text{CO}_2$ levels: 280 μatm (pre-industrial), 400 μatm (present-day) and 800 μatm (as projected for the end of this century).

2. MATERIAL AND METHODS

(a) Study sites and sampling

CWCs and ambient sea water were sampled during several cruises: June 2009 in the canyon of Lacaze-Duthiers ($42^\circ 32.98' \text{N}$; $03^\circ 25.21' \text{E}$), in September 2009 in the Gulf of Cassidaigne ($43^\circ 06.76' \text{N}$; $05^\circ 27.71' \text{E}$), both on board the R/V *Minibex* (COMEX, France), and in December 2009 in the Strait of Sicily at a site southwest off the island of Malta ($35^\circ 49.77' \text{N}$; $14^\circ 04.90' \text{E}$) on board the R/V *Urania* (Italy; figure 1).

During the *Minibex* cruises, CWCs were sampled with a remotely operated vehicle (ROV), while during the R/V *Urania* cruise a mini-dredge had to be deployed for a hit-and-go sampling because of technical problems with the ROV. At all sites, sea water next to living CWC was sampled using 6 l or 10 l Niskin bottles. After collection, corals were directly transferred to a plastic container ($1040 \times 640 \times 515 \text{ mm}$), containing about 120 l of sea water, which was continuously flowing through two layers of micron bags (5 and $1 \mu\text{m}$) and a chilling unit at a rate of 1000 l h^{-1} . Temperature was maintained at $12.5 \pm 0.5^\circ \text{C}$ (figure 2). Sea water was sub-sampled to determine the concentration of inorganic nutrients (phosphate and ammonium), total alkalinity (A_T) and dissolved inorganic carbon (C_T). Another




Figure 1. Sampling sites in the Mediterranean Sea: Lacaze-Duthiers (LD), Gulf of Cassidaigne (CS) and south of Malta (MC).




Figure 2. (a) Experimental set-up showing a large container with open and closed system incubation vials and some coral samples. In the right corner of the container is a double layer of micron bags (5 and $1 \mu\text{m}$) to filter the recycled and chilled sea water. Next to the container is the chiller unit that is connected to a pump (1000 l h^{-1}), which exchanges sea water between the container and chiller. (b) Aeration system to supply a multiple set of incubation vials. Tubing is joined by several silicon tubes with various diameters to fit to a standard aeration (4/6 mm inner/outer diameter) tube and an air pump. The space between tubing is sealed air-tight using silicone glue, and an (c) open system incubation vial containing a small branch of *M. oculata*. Inside the vial is another small tube of ca 1 cm in diameter in which the small 1 mm air tubing ends. The upward moving air bubbles inside the 1 cm tube generate an upward flow of sea water and thus water mixing.

part of the sea water collected from CWC sites was used for on-board incubation to determine calcification rates of corals by the alkalinity anomaly technique.

(b) On-board incubation to determine calcification rates of CWCs in ambient sea water

Small fragments were broken off from CWCs placed in either open or closed vials containing 150 ml or 180 ml sea water, respectively, and directly incubated after fragmentation (table 2 and figure 2). Sea water for incubation was sampled close to CWC sites to provide similar conditions with respect to the carbonate chemistry and nutrients. Open vials contained a small tube (\varnothing 1 cm) in which the outlet of a thin air tubing (\varnothing 3 mm) ended (figure 2b,c) to aerate the sea water with ambient air and to equilibrate the $p\text{CO}_2$ during incubation. In the open system set-up, the vials were aerated with ambient air, which served two purposes: (i) to keep $p\text{CO}_2$ stable and (ii) generate an air lift and thus a vertical flow and mixing of the incubation water. For closed incubation, vials were filled almost to the rim (180 ml) with sea water, and closed with a screw-on lid after corals were placed inside. Closed vials had a couple of millilitre of air space to avoid large changes in dissolved oxygen and to allow for some water mixing via ship movement. As controls, at least three vials contained no corals (blank) and also sea water was sub-sampled prior to starting incubation (T_0) to determine initial carbonate chemistry and inorganic nutrient concentration. Corals were incubated for 24 h, and then sea water was sub-sampled for determination of inorganic nutrients, C_T and A_T .

(c) Repeated incubation of CWCs under variable $p\text{CO}_2$

First, seven fragments were used in open system incubation and eight in closed system incubation to assess calcification rates using ambient sea water and thus ambient $p\text{CO}_2$ (table 2, CS5_500) directly on board. This on-board incubation served as a control to assess if calcification rates change over time (in a couple of days) and as a function of handling and transport. The same 15 coral fragments (seven open and eight closed system) were then used for repeated incubation at variable $p\text{CO}_2$ using the closed system set-up for all 15 coral fragments. About 2 h after transport back to the laboratory the first incubation was carried out at elevated $p\text{CO}_2$ (ca 867 μatm), then at ambient $p\text{CO}_2$ (404 μatm) and reduced $p\text{CO}_2$ (285 μatm) for 24 h each without feeding the corals. The timespan between the first on-board incubation and the last incubation at reduced $p\text{CO}_2$ was 5 days. Over this time period, we can also exclude that starvation affects calcification rates of CWCs [31]. For each treatment, at least three additional controls containing only sea water at respective $p\text{CO}_2$ (blank) were incubated along with coral samples. Except for ambient $p\text{CO}_2$, the $p\text{CO}_2$ level of bulk sea water was regulated before incubation using mass flow controllers (AnalytMTC, GFC 171) using a mix of CO_2 -free air and pure CO_2 . CO_2 -free air was produced by stripping CO_2 from ambient using soda lime. $p\text{CO}_2$ was measured using an infrared gas analyser (LI-COR-6262).

(d) Analyses of inorganic nutrients

Approximately 30 ml of sea water from Niskin bottles or experimental incubation vials were sub-sampled for inorganic nutrient analysis. Most of the sea water was used for rinsing syringe, filter and vials before a final volume of ca 5 ml of sea water was filtered over a 0.2 μm Acrodisc into 6 ml Pony Vials (Perkin Elmer). The samples were

frozen and stored at -20°C pending analysis at the Royal Netherlands Institute for Sea Research. Ammonium and dissolved inorganic phosphorus concentrations were determined using an AxFlow Bran & Luebbe Traacs800 autoanalyser [32]. Precision (± 1 s.d.) of an internal standard was 0.009 $\mu\text{mol kg}^{-1}$ and 0.023 $\mu\text{mol kg}^{-1}$ for phosphate and ammonium ($n = 17$ each), respectively.

(e) Analysis of A_T and C_T and determination of sea water carbonate chemistry

Ambient sea water was sampled to analyse the dissolved inorganic carbon (C_T) and total alkalinity (A_T) and to calculate other parameters of the carbonate system. Triplicate 500 ml samples were collected from a Niskin bottle. Sea water was poisoned with 100 μl saturated mercuric chloride, the stoppers of the bottles was greased (Apiezon) to seal bottles air-tight, and samples were stored at 10°C pending analysis. Analyses of C_T and A_T were performed by the Service National d'Analyse des Paramètres Océaniques du CO_2 (<http://soon.ipsl.jussieu.fr/SNAPCO2/>) following the method of Edmond [33] and the software described by DOE [34]. Reproducibility of the Dickson standard (batch 99, <http://www.andrew.uc.edu/co2qc/batches.html>) was 3 $\mu\text{mol kg}^{-1}$ for A_T and 2.8 $\mu\text{mol kg}^{-1}$ for C_T .

Initial and final C_T and A_T of the incubations were measured on 125 ml samples at the Laboratoire d'Océanographie de Villefranche. They were processed and stored as described earlier for the *in situ* samples. Bottles were carefully filled as described for samples from Niskin bottles, poisoned with 25 μl mercuric chloride, sealed air-tight and stored at 10°C .

C_T was determined immediately after opening the bottle using an inorganic carbon analyser (AIRICA, Marianda, Kiel, Germany) coupled to an infrared gas analyser (LI-COR 6262). This system was calibrated prior to sample analysis against a certified standard [34].

The sample bottle was kept in a water bath at a constant temperature (25°C). The coefficient of variation of at least three replicate measurements was 0.08 per cent and 2 $\mu\text{mol kg}^{-1}$. Also, an internal laboratory standard was measured at regular intervals and revealed a precision of $\pm 2.54 \mu\text{mol kg}^{-1}$ (mean = 2281.87, $n = 17$).

A_T was determined on samples filtered on 0.2 μm membranes by a potentiometric titration (Metrohm Titrand 80 titrator) coupled to a glass electrode (Metrohm, electrode plus) and thermometer. The pH electrode was calibrated on the total scale using Tris/HCl and 2-aminopyridine/HCl buffer solutions with a salinity of 38 (Dickson and co-workers [35]). Measurements were carried out in triplicate at 25°C with pre-weighed samples of ca 25 g and 0.1 N HCl (TitriPAC, Merck) was used for titration. A_T was calculated as described by Dickson and co-workers [35]. Standards provided by Dickson (batch 99 and 102) were run regularly. Accuracy was on average 2.38 $\mu\text{mol kg}^{-1}$ and 0.94 $\mu\text{mol kg}^{-1}$ below the nominal values of batches 99 and 102, respectively. Precision (± 1 s.d.) of repeated measurements was 3.53 and 4.63 $\mu\text{mol kg}^{-1}$ for the same batches ($n = 34$ and $n = 70$). Other parameters ($p\text{CO}_2$, Ω_a , pH) of the carbonate chemistry were determined from C_T , A_T , temperature, depth, and salinity using the software package SEACARB [36].

(f) Determination of calcification rates (G) by the alkalinity anomaly technique

The alkalinity anomaly technique was used to estimate rates of calcification using the 2 : 1 stoichiometric relationship between

Table 1. *In situ* sea water chemistry of CWC sampling sites. (Total alkalinity (A_T), concentration of dissolved inorganic carbon (C_T), bicarbonate (HCO_3^-), carbonate (CO_3^{2-}), phosphate (PO_4) and ammonium (NH_4) (all in $\mu\text{mol kg}^{-1}$). Also shown are pCO_2 (μatm), pH on total scale (pH_T), and the saturation state of aragonite (Ω_a). Numbers given in italics were calculated from T_A and C_T at respective depth (metre), salinity and temperature using the software package SEACARB, under the program R. For the stations of Lacaze-Duthiers (LD_9, LD_10) and the Gulf of Cassidaigne (CS_5) salinity and temperature were 38°C , and 13°C , and at the site south of Malta (MC_30) they were 38.75°C , and 12.75°C , respectively.)

station	depth	A_T	C_T	HCO_3^-	CO_3	CO_2	pCO_2	Ω_a	pH_T	PO_4	NH_4
LD_9	500	2577	2317	<i>2114</i>	<i>187</i>	<i>15.4</i>	<i>395</i>	2.6	<i>8.07</i>	0.41	0.29
LD_10	267	2577	2316	<i>2112</i>	<i>189</i>	<i>15.4</i>	<i>394</i>	2.7	<i>8.08</i>	0.22	0.16
CS_5	200	2615	2333	<i>2115</i>	<i>204</i>	<i>14.4</i>	<i>368</i>	3.0	<i>8.11</i>	0.28	0.20
MC_30	690	2623	2328	<i>2104</i>	<i>210</i>	<i>13.7</i>	<i>349</i>	2.8	<i>8.11</i>	0.20	0

Table 2. Results from on-board incubation of CWC in ambient sea water at three Mediterranean CWC sites: Lacaze-Duthiers (LD), Gulf of Cassidaigne (CS) and south of Malta (MC), numbers no._no. indicate station number and depth, respectively. (The three CWC species were *L. pertusa* (LP), *M. oculata* (MO) and *D. dianthus* (DE). Corals were either incubated in a closed or open system, DW is skeletal dry weight, average pCO_2 and Ω_a are average between beginning and end of incubation, G is calcification rate (corrected for inorganic nutrient excretion). CS5_500*: initially, seven corals were incubated, but only three could be evaluated with respect to carbonate chemistry and G in the open system. Nevertheless, DW and no. of polyps are given for all seven fragments because they were later used along with the eight fragments of the closed system of CS5_500 for repeated measurements of G under variable pCO_2 (figure 4).)

station	species	open/ closed	n	DW (G)	no. of polyps	initial pCO_2	average pCO_2	Ω_a	G ($\% \text{d}^{-1}$)
LD9_500	LP	open	3	0.49 ± 0.09	3.3 ± 1.5	354	378 ± 4	2.6 ± 0	0.090 ± 0.020
LD10_267	LP	open	3	2.03 ± 0.72	8.3 ± 1.5	354	389 ± 10	2.5 ± 0.1	0.033 ± 0.011
LD9_500	LP	closed	3	3.36 ± 2.63	7.0 ± 1.7	347	441 ± 56	2.1 ± 0.4	0.034 ± 0.013
LD9_500	MO	closed	3	2.21 ± 1.38	n.a. \pm n.a.	347	387 ± 20	2.5 ± 0.2	0.018 ± 0.018
CS5_500	MO	open	7	1.42 ± 0.15	47.1 ± 10.6	n.a.	n.a.	n.a.	n.a.
CS5_500*	MO	open	3	1.33 ± 0.10	40.3 ± 5.0	367	444 ± 127	2.4 ± 0.8	0.021 ± 0.017
CS5_500	MO	closed	8	1.03 ± 0.42	43.3 ± 12.5	404	863 ± 588	1.5 ± 0.8	0.068 ± 0.068
MC25_690	LP	open	1	3.53 ± 0	$7.0 \pm$ n.a.	433	$686 \pm$ n.a.	1.3 ± 0	0.028 ± 0
MC25_690	MO	open	6	2.79 ± 2.38	23.5 ± 8.3	433	614 ± 72	1.6 ± 0.3	0.037 ± 0.036
MC25_690	DE	open	5	4.49 ± 2.50	3.4 ± 1.1	433	563 ± 57	1.8 ± 0.2	0.011 ± 0.007
all groups			40	2.15 ± 1.85	25.4 ± 20.1	390	585 ± 340	1.9 ± 0.6	0.042 ± 0.040

the decrease of A_T and the precipitation of CaCO_3 [37]. However, other biogeochemical processes, such as changes in inorganic nutrient concentration can also affect A_T [35]. Previous work has shown that the CWC *L. pertusa* can release significant amounts of inorganic nutrient during incubation [38]. Changes of A_T during the incubations were therefore corrected for the release of phosphate and ammonium as follows:

$$G = -2\Delta T_A + \Delta \text{NH}_3 + \Delta \text{PO}_4.$$

Calcification rates were normalized to initial skeletal weight of the corals and are given in per cent per day ($\% \text{d}^{-1}$).

(g) Statistical analyses

Statistical analyses were conducted using the software package STATISTICA v. 7.0. The tests used are mentioned in the results subsections. Values are reported as mean \pm standard deviation.

For parameters of the carbonate chemistry and inorganic nutrients at the start and end of the experiments, see tables in the electronic supplementary material.

3. RESULTS

(a) *In situ* sea water carbonate chemistry and inorganic nutrients

A_T and C_T of ambient sea water ranged from 2577 to 2623 and 2316 to 2333, respectively (table 1). The calculated pCO_2 , pH_T and Ω_a values ranged between

349 μatm and 395 μatm , 8.07 and 8.11, and 2.58 and 2.95, respectively. Average phosphate and ammonium concentrations ranged between 0.20 $\mu\text{mol kg}^{-1}$ and 0.41 $\mu\text{mol kg}^{-1}$ and 0.03–0.29 $\mu\text{mol kg}^{-1}$, respectively.

(b) Changes of inorganic nutrient concentrations during the incubations

The average phosphate and ammonium release during the incubations was $0.06 \pm 0.07 \mu\text{mol g}^{-1}$ and $1.12 \pm 1.19 \mu\text{mol g}^{-1}$ skeleton per day, respectively. The lowest release of phosphate was found for *Desmophyllum dianthus* with $0.002 \pm 0.002 \mu\text{mol g}^{-1}$ skeleton per day ($n = 2$), while in *M. oculata* and *L. pertusa* incubations, phosphate release was $0.06 \pm 0.07 \mu\text{mol g}^{-1}$ skeleton per day ($n = 27$) and $0.08 \pm 0.06 \mu\text{mol g}^{-1}$ skeleton per day ($n = 10$), respectively. For ammonium, the highest release rates were found for *M. oculata* with $1.32 \pm 1.37 \mu\text{mol g}^{-1}$ skeleton per day, while *L. pertusa* released an average of $0.77 \pm 0.53 \mu\text{mol g}^{-1}$ skeleton per day and *D. dianthus* $0.92 \pm 0.53 \mu\text{mol g}^{-1}$ skeleton per day. Between species effects were tested for *L. pertusa* and *M. oculata*, while inorganic nutrient release was only determined for two *D. dianthus* specimens and was thus not considered for statistical testing. There was no species-specific effect on the release of phosphate or ammonium (t -test, $t = 0.65$, $p = 0.52$ for phosphate and $t = -1.22$, $p = 0.23$ for ammonium, $n = 35$ for both nutrients). Also, phosphate




Figure 3. Correlation between G corrected for inorganic nutrient increase during incubation and uncorrected ($r^2 = 0.991$, $p \ll 0.001$; $n = 37$).




Figure 4. Average calcification rates of *M. oculata* of repeated closed system incubation at high (867 μatm), ambient (445 μatm) and low (285 μatm) pCO_2 . Data are from repeated incubations in the laboratory shortly after return from the cruise using the same coral fragments ($n = 15$). A repeated-measures ANOVA revealed a significant effect of pCO_2 on calcification. Vertical bars denote 0.95 confidence intervals.

and ammonium excretion exhibit a positive and significant correlation when normalized to skeletal weight ($R = 0.79$, $p \ll 0.001$, $n = 39$).

(c) Contribution of changes in nutrient concentration to changes in A_T

There is a significant correlation between the release of nutrients and the rate of calcification (PO_4 : $R = 0.72$, NH_4 : $R = 0.79$, $p \ll 0.001$ for both, $n = 39$). The calcification rates estimated without (G_{uncorr}) and with (G) correction for changes in the concentration of nutrients are significantly correlated (figure 3; $G = 1.11 \times G_{\text{uncorr}} + 0.001$, $r = 0.995$, $p \ll 0.001$; $n = 37$). Therefore, the rate of calcification is underestimated by ca 10 per cent if nutrient excretion is not accounted for. As not all incubation vials were sub-sampled for inorganic nutrients, this linear regression function was used to correct for nutrient excretion where respective data were missing.

(d) Calcification rates in ambient sea water

The overall mean calcification rates of corals incubated in ambient sea water was $0.04 \pm 0.04\% \text{ d}^{-1}$ and ranged

between -0.01 and $0.23\% \text{ d}^{-1}$ ($n = 36$). Species-specific rates for *L. pertusa*, *M. oculata* and *D. dianthus* were $0.05 \pm 0.03\% \text{ d}^{-1}$ ($n = 10$), $0.05 \pm 0.05\% \text{ d}^{-1}$ ($n = 21$) and $0.01 \pm 0.01\% \text{ d}^{-1}$ ($n = 5$). Rates of calcification of *L. pertusa* were on average higher in the open incubation system with $0.05 \pm 0.04\% \text{ d}^{-1}$ ($n = 7$) versus $0.03 \pm 0.01\% \text{ d}^{-1}$ ($n = 3$) in the closed system, while the opposite was true for *M. oculata* with $0.03 \pm 0.03\% \text{ d}^{-1}$ ($n = 9$) and $0.06 \pm 0.06\% \text{ d}^{-1}$ ($n = 12$) in open and closed systems, respectively (table 2). There was no significant species effect on calcification rates between *M. oculata* and *L. pertusa*, between open and closed system incubation nor a combined effect of species and type of incubation (two-way ANOVA, $p = 0.98$ for the species effect, $p = 0.93$ for the incubation effect and $p = 0.23$ for the interaction term).

(e) Repeated incubations of *Madrepora oculata* under various pCO_2 levels

During the on-board incubation, there was a problem with the open system and only three of the seven corals were sub-sampled for A_T . Calcification of the three corals was $0.02 \pm 0.02\% \text{ d}^{-1}$ and $0.03 \pm 0.03\% \text{ d}^{-1}$ for on-board and laboratory (closed system) incubation, respectively. Calcification rates for the corals incubated in the closed incubation system were $0.07 \pm 0.07\% \text{ d}^{-1}$ for both on-board and laboratory incubation. Under ambient pCO_2 , paired t -tests revealed no significant difference between on-board and laboratory incubation (open system: $p = 0.74$, $n = 3$; closed system: $p = 0.93$, $n = 8$). Pooling the calcification data for the 11 corals (open and closed system), average calcification rates were 0.06 ± 0.06 and $0.06 \pm 0.07\% \text{ d}^{-1}$ and were not significantly different from each other (paired t -test: $p = 0.84$, $n = 11$).

The initial pCO_2 levels for the three subsequent experiments were 867, 445 and 285 μatm , corresponding to Ω_a values of 1.9, 2.6 and 3.7, respectively. pCO_2 increased during the closed system incubations owing to respiration and calcification. The average final pCO_2 levels were 1186 ± 461 , 629 ± 248 and $331 \pm 57 \mu\text{atm}$, which decreased Ω_a to 1.3 ± 0.6 , 1.8 ± 0.6 and 2.9 ± 0.6 , respectively.

Average calcification rates were 0.06 ± 0.06 , 0.06 ± 0.06 and $0.12 \pm 0.06\% \text{ d}^{-1}$ for high, ambient and low pCO_2 levels, respectively (figure 4). Repeated-measures ANOVA revealed that pCO_2 had a significant effect on calcification rates ($F_{2,28} = 11.3$, $p < 0.001$). *Post hoc* comparison showed, that G of high and ambient pCO_2 levels were significantly different from the low pCO_2 (Tukey honest significant difference test (HSD): $p = 0.001$ and $p < 0.001$, respectively), while G at ambient and high pCO_2 were not significantly different (Tukey HSD: $p = 0.98$).

4. DISCUSSION

(a) Methodological constraints

(i) On-board experiments

Measuring calcification of CWCs on board allows investigation of corals in ambient sea water over a broad biogeographic range. It requires a method that is not logistically demanding in terms of laboratory facilities. The set-up used for maintaining and incubating corals served that purpose well. It uses small incubation chambers that enable measurement of changes in total alkalinity large enough to estimate rates of calcification with satisfactory

replication. We also tested the use of a closed incubation system. In closed systems, the CO₂ released to the incubation water by coral respiration and calcification changes the sea water carbonate chemistry during incubation. This is likely to affect calcification rates as pCO₂ increases and Ω_a and pH decrease. However, systematic comparison revealed no significant difference in calcification between open and closed system incubation. Working with relatively simple and small incubation systems makes replication easier. This is an important experimental prerequisite given the high natural growth variability of CWCs that is largely owing to the age composition of polyps of a branch [16,21] and sporadic skeletal growth [16,18].

(ii) Pitfalls of alkalinity anomaly technique

The three most common methods to determine calcification rates of corals are ⁴⁵Ca-labelling [39], the buoyant weight determination [40] and the A_T anomaly technique [37]. The alkalinity anomaly technique used in this study has the advantage that it is sensitive enough to use for short-term incubations and that corals can be kept alive. However, it has been shown that *L. pertusa* and *M. oculata* excrete inorganic nutrients and that bacterial growth is stimulated during incubations owing to the release of organic matter [38,41,42]. This may also alter the T_A during incubation. Negatively charged surface groups of phytoplankton and bacteria can react during titration with hydrochloric acid and the acid-base functional groups of dissolved organic carbon (DOC) in sea water can also have a small influence on T_A [43]. It is therefore critical to pre-filter sea water over a 0.2 μ m Acrodisc. Changes in the DOC pool during incubation was not accounted for in the present study, but Kim *et al.* [43] suggested that the influence of DOC on A_T might be rather small. The release of inorganic nutrients induces a 10 per cent underestimation of calcification which was accounted for by an empirical relationship (this study). Also, a 10 per cent bias is relatively small when compared with the large natural variability of CWC calcification, with a coefficient of variation of up to 100 per cent.

(b) Calcification under ambient conditions

Calcification data are now available on *L. pertusa* determined by the three methods mentioned earlier. Calcification ranges between 0.02 and 0.07% d⁻¹ under initial ambient conditions for corals collected in Mingulay and Skagerrak and investigated with ⁴⁵Ca [21]. For Mediterranean samples, Orejas *et al.* [20] estimated calcification rates of 0.02% d⁻¹ using the buoyant weight technique, and rates ranging between 0.03 and 0.09% d⁻¹ were found in the present study with the alkalinity anomaly technique. The rate of calcification of the Mediterranean species *D. dianthus* and *M. oculata* determined by buoyant weight were 0.04% d⁻¹ and 0.11% d⁻¹, respectively [20], and 0.01 and 0.02–0.07% d⁻¹ determined by the alkalinity anomaly technique (this study). Interestingly, there seems to be no systematic bias to either higher or lower calcification rates between the methods used, which provides some confidence that all three methods provide similar estimates. The method used by Orejas *et al.* [20] was based on long-term aquarium maintenance (256 days), while the ⁴⁵Ca and A_T approaches were based on short-term on-board incubations (1 day) with freshly collected colonies. Our results suggest that calcification of *L. pertusa* and *M. oculata* is similar. By contrast, Orejas *et al.* [20] reported a

calcification rate fivefold higher in *M. oculata* than in *L. pertusa* and they hypothesize that this might be owing to different allocation of energy into tissue and skeletal growth. However, there is no evidence yet that *M. oculata* allocates more energy into skeletal growth than *L. pertusa* or *D. dianthus*.

(i) Size matters

It has been demonstrated in *L. pertusa* that calcification is four orders of magnitude lower in older than in younger polyps [38]. Hence, calcification rates normalized to skeletal weight diminish relative to increasing size and age of a colony. There is no information available on the weight of coral fragments used in the study of Orejas *et al.* [20], but the average polyp numbers were 2.9 and 6.2 polyps per fragment for *L. pertusa* and *M. oculata*, respectively. This means these fragments were very small with half and one-fifth of the polyp number used in the present study, respectively. One exception was the size of *L. pertusa* used in the first open system incubation (table 2) of the present study, as these specimens were as small as the ones used by Orejas *et al.* [20]. The calcification rate of these small fragments was 0.09% d⁻¹ while a rate of 0.02% d⁻¹ was found by Orejas *et al.* [20]. This indicates that the long-term aquarium conditions were less favourable for calcification of *L. pertusa* of the same size class than were the short-term on-board incubations, while the calcification rates of *M. oculata* reported by the two studies are in good agreement.

(ii) Site-specific calcification rates

The present study also reports calcification measured in three distinct sites, providing the opportunity to assess geographical differences. Although, the Mediterranean exhibits an east–west trend in nutrient concentration [44] and in anthropogenic CO₂ uptake [29], the *in situ* carbonate chemistry was very similar between the sites (table 1) and there is no significant difference in calcification rate between sites. However, on a broader biogeographic and depth scale where parameters of the carbonate chemistry are more distinct from each other, patterns in calcification rates might exist. *In situ* studies would be the ideal method to investigate the response of CWCs to changing environmental conditions, but are logistically too demanding for empirical comparisons on a broad geographical and depth scale. However, it is important to gather comparable and empirical datasets as close to *in situ* as possible to derive a baseline understanding, and moreover, to better predict the potential response to environmental change scenarios.

(c) Response to reduced and elevated pCO₂

Our results clearly show that there is no negative response on calcification rates at the higher pCO₂ level (867 μ atm). However, calcification almost doubled at the pre-industrial pCO₂ level (285 μ atm) relative to calcification under the control at ambient pCO₂. So far, only one study has tested the effect of ocean acidification on the calcification of CWCs. It revealed a negative response with decreasing pH (higher pCO₂) in *L. pertusa* from the Skagerrak [21]. However, in contrast to the study on *L. pertusa*, Ω_a did not drop below 1 for *M. oculata* in the higher pCO₂ treatment, which might explain why *M. oculata* did not reveal a similar negative response at higher pCO₂. While for most reef-building zooxanthellate corals a negative and often linear response to increasing pCO₂ levels and lower Ω_a had been reported [45–47],

it had already been shown for temperate corals that calcification rates can remain unaffected by pCO₂ levels as high as that projected for the end of the century [48,49]. In the study of Ries *et al.* [48], a nonlinear response was shown with calcification rates of *Oculina arbuscula* remaining constant at pCO₂ levels between 400 and 900 µatm and decreasing at a pCO₂ of 2850 µatm. It is therefore possible that *M. oculata* exhibits a similar nonlinear response and additional studies would be necessary to determine at which pCO₂ level (or Ω_a) calcification rates start to decrease. Few studies are actually available where calcification rates were tested for lower pCO₂ levels. For the tropical coral *Porites lutea*, a linear response to Ω_a, and increasing calcification rates with decreasing pCO₂ were revealed [50]. For the Mediterranean Sea, it has been shown, that pH has decreased by 0.1 pH units owing to anthropogenic CO₂ [29] and our results thus indicate that ocean acidification already has had a detrimental effect on calcification rates of *M. oculata* with a reduction by 50 per cent since pre-industrial time. *Desmophyllum dianthus*, *M. oculata* and *L. pertusa* have distinct sensitivities to increased temperature, with *L. pertusa* being the least tolerant to higher temperatures (C. Maier 2009, personal observation). This species is probably already at its highest temperature limit in the Mediterranean Sea. If the pCO₂ increase since pre-industrial time had a similarly negative effect on *L. pertusa* calcification than it had on *M. oculata*, it could mean that *L. pertusa* might already be more negatively affected by climate change in the Mediterranean, where both temperature rise [26] and anthropogenic CO₂ [29] are well documented. This might also partly explain, why *M. oculata* is, at the present-day, more widespread in the Mediterranean Sea than *L. pertusa* despite the fact that in the North Atlantic *L. pertusa* is the dominant CWC species. However, more studies including other parameters that may influence coral growth are needed to draw a final conclusion on a species-specific response to climate change scenarios.

We thank captains, crew and scientific shipboard staff of R/V *Minibex*, and R/V *Urania*, respectively. Cruise MEDCOR was partially funded by CNR and the EU HERMES programme, cruises on RV *Minibex* were part of a project on canyons of the Mediterranean Sea (MedSeaCan) lead by the Agences des aires marines protégées. Special thanks to L. Angeletti and A. Ceregato for on-board assistance, to S. Comeau for his assistance with the measurement of total alkalinity and J. Ries and an anonymous reviewer for their helpful comments on the manuscript. Financial support was provided by the European Commission through a Marie-Curie Fellowship (MECCA, project no 220299 and the project COMP via the Prince Albert II of Monaco Foundation. This work is a contribution to the 'European Project on Ocean Acidification' (EPOCA) which received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 211384, and also contributes to EU HERMIONE project (contract no. 226354) and is ISMAR-CNR Bologna scientific contribution 1733.

REFERENCES

- 1 Roberts, J. M., Wheeler, A. J. & Freiwald, A. 2006 Reefs of the deep: the biology and geology of cold-water coral ecosystems. *Science* **312**, 543–547. (doi:10.1126/science.1119861)
- 2 Freiwald, A., Beuck, L., Rüggeberg, A., Taviani, M. & Hebbeln, D. 2009 The white coral community in the central Mediterranean Sea revealed by ROV surveys. *Oceanography* **22**, 36–52. (doi:10.5670/oceanog.2009.06)
- 3 Heifetz, J. 2002 Coral in Alaska: distribution, abundance, and species associations. *Hydrobiologia* **471**, 19–28. (doi:10.1023/A:1016528631593)
- 4 Hovland, M., Vasshus, S., Indreeide, A., Austdal, L. & Nilsen, Ø. 2002 Mapping and imaging deep-sea coral reefs off Norway, 1982–2000. *Hydrobiologia* **471**, 13–17. (doi:10.1023/A:1016576514754)
- 5 Orejas, C., Gori, A., Iacono, C. L., Puig, P., Gili, J.-M. & Dale, M. R. T. 2009 Cold-water corals in the Cap de Creus canyon, northwestern Mediterranean: spatial distribution, density and anthropogenic impact. *Mar. Ecol. Prog. Ser.* **397**, 37–51. (doi:10.3354/meps08314)
- 6 Roberts, J. M., Brown, C. J., Long, D. & Bates, C. R. 2005 Acoustic mapping using a multibeam echosounder reveals cold-water coral reefs and surrounding habitats. *Coral Reefs* **24**, 654–669. (doi:10.1007/s00338-005-0049-6)
- 7 Taviani, M., Freiwald, A. & Zibrowius, H. 2005 Deep coral growth in the Mediterranean Sea: an overview. In *Cold-water corals and ecosystems* (eds A. Freiwald & J. M. Roberts), pp. 137–156. Heidelberg, Germany: Springer.
- 8 Taviani, M. *et al.* 2010 Pleistocene to Recent scleractinian deep-water corals and coral facies in the eastern Mediterranean. *Facies* **57**, 579–603. (doi:10.1007/s10347-010-0247-8)
- 9 White, M., Mohn, C., Stigter, H. & Mottram, G. 2005 Deep-water coral development as a function of hydrodynamics and surface productivity around the submarine banks of the Rockall Trough, NE Atlantic. In *Cold-water corals and ecosystems* (eds A. Freiwald & J. M. Roberts), pp. 503–514. Heidelberg, Germany: Springer.
- 10 Freiwald, A., Henrich, R. & Pätzold, J. 1997 Anatomy of a deep-water coral reef mound from Stjærnsund, West-Finnmark, Northern Norway. *SEPM Spec. Vol.* **56**, 141–162.
- 11 Mikkelsen, N., Erlenkauser, H., Killingley, J. S. & Berger, W. H. 1982 Norwegian corals: radiocarbon and stable isotopes in *Lophelia pertusa*. *Boreas* **5**, 163–171.
- 12 Mortensen, P. B. & Rapp, H. T. 1998 Oxygen and carbon isotope ratios related to growth line patterns in skeletons of *Lophelia pertusa* (L.) (Anthozoa, Scleractinia): implications for determination of linear extension rates. *Sarsia* **83**, 443–446.
- 13 Rogers, A. D. 1999 The biology of *Lophelia pertusa* (Linnaeus 1758) and other deep-water reef-forming corals and impacts from human activities. *Internat. Rev. Hydrobiol.* **84**, 315–406.
- 14 Bell, N. & Smith, J. 1999 Coral growing on North Sea oil rigs. *Nature* **402**, 601. (doi:10.1038/45127)
- 15 Gass, S. E. & Roberts, J. M. 2006 The occurrence of the cold-water coral *Lophelia pertusa* (Scleractinia) on oil and gas platforms in the North Sea: colony growth, recruitment and environmental controls on distribution. *Mar. Poll. Bull.* **52**, 549–559. (doi:10.1016/j.marpolbul.2005.10.002)
- 16 Mortensen, P. B. 2001 Aquarium observations on the deep-water coral *Lophelia pertusa* (L., 1758) (Scleractinia) and selected associated invertebrates. *Ophelia* **54**, 83–104.
- 17 Orejas, C., Gori, A. & Gili, J. M. 2008 Growth rates of live *Lophelia pertusa* and *Madrepora oculata* from the Mediterranean Sea maintained in aquaria. *Coral Reefs* **27**, 255. (doi:10.1007/s00338-007-0350-7)
- 18 Brooke, S. & Young, C. M. 2009 *In situ* measurement of survival and growth of *Lophelia pertusa* in the northern Gulf of Mexico. *Mar. Ecol. Prog. Ser.* **397**, 153–161. (doi:10.3354/meps08344)
- 19 Buddemeier, R. W. 1978 Coral growth: retrospective analysis. In *Coral reefs: research methods* (eds D. R. Stoddart & R. E. Johannes), pp. 551–571. Paris, France: Unesco.

- 20 Orejas, C., Ferrier-Pagès, C., Reynaud, S., Gori, A., Beraud, E., Tsounis, G., Allemand, D. & Gili, J. M. 2011 Long-term growth rates of four Mediterranean cold-water coral species maintained in aquaria. *Mar. Ecol. Prog. Ser.* **429**, 57–65. (doi:10.3354/meps09104)
- 21 Maier, C., Hegeman, J., Weinbauer, M. G. & Gattuso, P. 2009 Calcification of the cold-water coral *Lophelia pertusa* under ambient and reduced pH. *Biogeosciences* **6**, 1671–1680. (doi:10.5194/bg-6-1671-2009)
- 22 Guinotte, J. M., Orr, J., Cairns, S., Freiwald, A., Morgan, L. & George, R. 2006 Will human-induced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals? *Front Ecol. Environ.* **4**, 141–146. (doi:10.1890/1540-9295(2006)004[0141:WHCISC]2.0.CO;2)
- 23 Orr, J. C. et al. 2005 Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* **437**, 681–686. (doi:10.1038/nature04095)
- 24 Turley, C. M., Roberts, J. M. & Guinotte, J. M. 2007 Corals in deep-water: will the unseen hand of ocean acidification destroy cold-water ecosystems? *Coral Reefs* **26**, 445–448. (doi:10.1007/s00338-007-0247-5)
- 25 Bethoux, J. P., Gentili, B., Morin, P., Nicolas, E., Pierre, C. & Ruiz-Pino, D. 1999 The Mediterranean Sea: a miniature ocean for climatic and environmental studies and a key for the climatic functioning of the North Atlantic. *Prog. Oceanogr.* **44**, 131–146. (doi:10.1016/S0079-6611(99)00023-3)
- 26 Bethoux, J. P., Gentili, B., Raunet, J. & Tailliez, D. 1990 Warming trend in the western Mediterranean deep water. *Nature* **347**, 660–662. (doi:10.1038/347660a0)
- 27 Lelieveld, J., Berresheim, H., Borrmann, S. & Crutzen, P. J. 2002 Global air pollution crossroads over the Mediterranean. *Science* **298**, 794–799. (doi:10.1126/science.1075457)
- 28 Nykjaer, L. 2009 Mediterranean Sea surface warming 1985–2006. *Clim. Res.* **39**, 11–17. (doi:10.3354/cr00794)
- 29 Touratier, F. & Goyet, C. 2011 Impact of the eastern Mediterranean transient on the distribution of anthropogenic CO₂ and first estimate of acidification for the Mediterranean Sea. *Deep Sea Res. I* **58**, 1–15.
- 30 CIESM 2008 Impacts of ocean acidification on biological, chemical and physical systems in the Mediterranean and Black Seas. In *CIESM Workshop Monographs*, vol. 36 (ed. F. Briand), pp. 124. Monaco.
- 31 Schubert, A. 2010 *The impact of ocean acidification on calcification of Mediterranean cold-water corals*. Diplomarbeit, IFM-GEOMAR, pp. 68. Germany: Christian Albrechts-Universität zu Kiel.
- 32 Herfort, L., Schouten, S., Abbas, B., Veldhuis, M. J. W., Coolen, M. J. L., Wuchter, C., Boon, J. P., Herndl, G. J. & Damsté, J. S. S. 2007 Variations in spatial and temporal distribution of Archaea in the North Sea in relation to environmental variables. *FEMS Microbiol. Ecol.* **62**, 242–257. (doi:10.1111/j.1574-6941.2007.00397.x)
- 33 Edmond, J. M. 1970 High precision determination of titration alkalinity and total carbon dioxide content of seawater by potentiometric titration. *Deep Sea Res.* **17**, 737–750.
- 34 DOE 1994 Recommended standard operation procedures. In *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water*, version 2 (eds A. G. Dickson & C. Goyet), pp. 1–187. (ORNL/CDIAC-74). Online version.
- 35 Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Körtzinger, A. & Dickson, A. G. 2007 Total alkalinity: the explicit conservative expression and its application to biogeochemical processes. *Deep Sea Res. I* **106**, 287–300.
- 36 Lavigne, H. & Gattuso, P. 2011 SEACARB: seawater carbonate chemistry with R. R package version 2.4. See <http://www.CRAN.R-project.org/package=seacarb>.
- 37 Chisholm, J. R. M. & Gattuso, P. 1991 Validation of the alkalinity anomaly technique for investigating calcification and photosynthesis in coral reef communities. *Limnol. Oceanogr.* **36**, 1232–1239. (doi:10.4319/lo.1991.36.6.1232)
- 38 Maier, C., de Kluijver, A., Agis, M., Brussaard, C. P. D., Van Duyl, F. C. & Weinbauer, M. G. 2011 Dynamics of nutrients, total organic carbon, prokaryotes and viruses in onboard incubations of cold-water corals. *Biogeosci. Discuss.* **8**, 3829–3861. (doi:10.5194/bgd-8-3829-2011)
- 39 Tambutté, É., Allemand, D., Mueller, E. & Jaubert, J. 1996 A compartmental approach to the mechanism of calcification in hermatypic corals. *J. Exp. Biol.* **199**, 1029–1041.
- 40 Davies, P. S. 1989 Short-term growth measurements of corals using an accurate buoyant weighing technique. *Mar. Biol.* **101**, 389–395. (doi:10.1007/BF00428135)
- 41 Wild, C., Mayr, C., Wehrmann, L., Schöttner, S., Naumann, M., Hoffmann, F. & Rapp, H. T. 2008 Organic matter release by cold-water corals and its implication for fauna-microbe interaction. *Mar. Ecol. Prog. Ser.* **372**, 67–75. (doi:10.3354/meps07724)
- 42 Wild, C., Wehrmann, L. M., Mayr, C., Schöttner, S. I., Allers, E. & Lundälv, T. 2009 Microbial degradation of cold-water coral-derived organic matter: potential implication for organic C cycling in the water column above Tisler Reef. *Aquat. Biol.* **7**, 71–80. (doi:10.3354/ab00185)
- 43 Kim, H.-C., Lee, K. & Choi, W. 2006 Contribution of phytoplankton and bacterial cells to the measured alkalinity of seawater. *Limnol. Oceanogr.* **51**, 331–338. (doi:10.4319/lo.2006.51.1.0331)
- 44 Bethoux, J. P., Morin, P., Chamuery, C., Connan, O., Gentili, B. & Ruiz-Pino, D. 1998 Nutrients in the Mediterranean Sea, mass balance and statistical analysis of concentrations with respect to environmental change. *Mar. Chem.* **63**, 155–169. (doi:10.1016/S0304-4203(98)00059-0)
- 45 Langdon, C. & Atkinson, M. J. 2005 Effect of elevated pCO₂ on photosynthesis and calcification of corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment. *J. Geophys. Res.* **110**, C09S07. (doi:10.1029/2004JC002576)
- 46 Marubini, F., Ferrier-Pagès, C., Furla, P. & Allemand, D. 2008 Coral calcification responds to seawater acidification: a working hypothesis towards a physiological mechanism. *Coral Reefs* **27**, 491–499. (doi:10.1007/s00338-008-0375-6)
- 47 Schneider, K. & Erez, J. 2006 The effect of carbonate chemistry on calcification and photosynthesis in the hermatypic coral *Acropora eurystroma*. *Limnol. Oceanogr.* **51**, 1284–1293. (doi:10.4319/lo.2006.51.3.1284)
- 48 Ries, J. B., Cohen, A. L. & McCorkle, D. C. 2010 A non-linear calcification response to CO₂-induced ocean acidification by the coral *Oculina arbuscula*. *Coral Reefs* **29**, 661–674. (doi:10.1007/s00338-010-0632-3)
- 49 Rodolfo-Metalpa, R., Martin, S., Ferrier-Pagès, C. & Gattuso, P. 2010 Response of the temperate coral *Cladocora caespitosa* to mid- and long-term exposure to pCO₂ and temperature levels projected for the year 2100 AD. *Biogeosciences* **7**, 289–300. (doi:10.5194/bg-7-289-2010)
- 50 Ohde, S. & Hossain, M. M. M. 2004 Effect of CaCO₃ (aragonite) saturation state of seawater on calcification of *Porites* coral. *Geochem. J.* **38**, 613–621. (doi:10.2343/geochemj.38.613)