

screen: the light by which the eye then sees the screen is, of course, undulatory.

The phenomena in these exhausted tubes reveal to physical science a new world—a world where matter exists in a fourth state, where the corpuscular theory of light holds good, and where light does not always move in a straight line; but where we can never enter, and in which we must be content to observe and experiment from the outside.

II. "On a Machine for the Solution of Simultaneous Linear Equations." By Sir WILLIAM THOMSON, LL.D., F.R.S., President of the Royal Society, Edinburgh. Received August 30, 1878.

Let  $B_1, B_2, \dots, B_n$  be  $n$  bodies each supported on a fixed axis (in practice each is to be supported on knife-edges like the beam of a balance).

Let  $P_{11}, P_{21}, P_{31}, \dots, P_{n1}$ ,  $n$  pulleys each pivoted on  $B_1$ ;  
 $P_{12}, P_{22}, P_{32}, \dots, P_{n2}$                    "                   "                    $B_2$ ;  
 $P_{13}, P_{23}, P_{33}, \dots, P_{n3}$                    "                   "                    $B_3$ ;

$C_1, C_2, C_3, \dots, C_n$ ,  $n$  cords passing over the pulleys;

$D_1, P_{11}, P_{12}, P_{13}, \dots, P_{1n}, E_1$ , the course of  $C_1$ ;

$D_2, P_{21}, P_{22}, P_{23}, \dots, P_{2n}, E_2$ ,                   "                    $C_2$ ;

$D_1, E_1, D_2, E_2, \dots, D_n, E_n$ , fixed points;

$l_1, l_2, l_3, \dots, l_n$  the lengths of the cords between  $D_1, E_1$ , and  $D_2, E_2, \dots$  and  $D_n, E_n$ , along the courses stated above, when  $B_1, B_2, \dots, B_n$  are in particular positions which will be called their zero positions;

$l_1 + e_1, l_2 + e_2, \dots, l_n + e_n$ , their lengths between the same fixed points, when  $B_1, B_2, \dots, B_n$  are turned through angles  $x_1, x_2, \dots, x_n$  from their zero positions;

(11), (12), (13),  $\dots$  (1*n*),

(21), (22), (23),  $\dots$  (2*n*),

(31), (32), (33),  $\dots$  (3*n*),

quantities such that

$$\left. \begin{aligned} (11)x_1 + (12)x_2 + \dots + (1n)x_n &= e_1 \\ (21)x_1 + (22)x_2 + \dots + (2n)x_n &= e_2 \\ (31)x_1 + (32)x_2 + \dots + (3n)x_n &= e_3 \\ \dots & \dots \\ (n1)x_1 + (n2)x_2 + \dots + (nn)x_n &= e_n \end{aligned} \right\} \dots \quad (I)$$

We shall suppose  $x_1, x_2, \dots, x_n$  to be each so small that (11), (12),





removed, and it is advisable to make the tensions each equal to half the weight of one of the pulleys with its adjustable frame.) The machine is now ready for use. To use it, pull the cords simultaneously or successively till lengths equal to  $e_1, e_2, \dots e_n$  are passed through the rings  $E_1, E_2, \dots E_n$ , respectively.

The *pulls* required to do this may be positive or negative; in practice, they will be infinitesimal, downward or upward pressures applied by hand to the stretching weights which (§) remain permanently hanging on the cords.

Observe the angles through which the bodies  $B_1, B_2, \dots B_n$  are turned by this given movement of the cords. These angles are the required values of the unknown  $x_1, x_2, \dots x_n$ , satisfying the simultaneous equations (I).

The actual construction of a practically useful machine for calculating as many as eight or ten or more of unknowns from the same number of linear equations does not promise to be either difficult or over-elaborate. A fair approximation being found by a first application of the machine, a very moderate amount of straightforward arithmetical work (aided very advantageously by Crelle's multiplication tables) suffices to calculate the residual errors, and allow the machines (with the setting of the pulleys unchanged) to be re-applied to calculate the corrections (which may be treated decimally, for convenience): thus, 100 times the amount of the correction on each of the original unknowns, to be made the new unknowns, if the magnitudes thus falling to be dealt with are convenient for the machine. There is, of course, no limit to the accuracy thus obtainable by successive approximations. The exceeding easiness of each application of the machine promises well for its real usefulness, whether for cases in which a single application suffices, or for others in which the requisite accuracy is reached after two, three, or more of successive approximations.

December 12, 1878.

W. SPOTTISWOODE, M.A., D.C.L., President, in the Chair.

Dr. Philipp Hermann Sprengel was admitted into the Society.

The Presents received were laid on the table, and thanks ordered for them.

The following Papers were read:—