

## Introduction



**Cite this article:** Westra ER, van Houte S, Gandon S, Whitaker R. 2019 The ecology and evolution of microbial CRISPR-Cas adaptive immune systems. *Phil. Trans. R. Soc. B* **374**: 20190101.  
<http://dx.doi.org/10.1098/rstb.2019.0101>

Accepted: 18 February 2019

One contribution of 17 to a discussion meeting issue 'The ecology and evolution of prokaryotic CRISPR-Cas adaptive immune systems'.

### Subject Areas:

ecology, evolution, genetics, microbiology

### Authors for correspondence:

Edze R. Westra

e-mail: [e.r.westra@exeter.ac.uk](mailto:e.r.westra@exeter.ac.uk)

Stineke van Houte

e-mail: [c.van-houte@exeter.ac.uk](mailto:c.van-houte@exeter.ac.uk)

Sylvain Gandon

e-mail: [sylvain.gandon@cefe.cnrs.fr](mailto:sylvain.gandon@cefe.cnrs.fr)

Rachel Whitaker

e-mail: [rwhitaker@life.illinois.edu](mailto:rwhitaker@life.illinois.edu)

# The ecology and evolution of microbial CRISPR-Cas adaptive immune systems

Edze R. Westra<sup>1</sup>, Stineke van Houte<sup>1</sup>, Sylvain Gandon<sup>2</sup> and Rachel Whitaker<sup>3</sup>

<sup>1</sup>ESI and CEC, Biosciences, University of Exeter, Cornwall Campus, Penryn TR10 9EZ, UK

<sup>2</sup>CEFE UMR 5175, CNRS Université de Montpellier Université Paul-Valéry Montpellier EPHE, 34293 Montpellier Cedex 5, France

<sup>3</sup>Department of Microbiology, University of Illinois, Urbana-Champaign, 601 S. Goodwin Avenue, Urbana, IL 61801, USA

ERW, 0000-0003-4396-0354; SG, 0000-0003-2624-7856; RW, 0000-0003-2263-5798

## 1. Introduction

Over the past decade, the field of CRISPR-Cas research has received a lot of attention from the scientific community. While initially, this mostly concerned microbiologists who were fascinated by the discovery that some bacteria encode RNA-guided adaptive immune systems, this rapidly spread to other scientific disciplines following the development of groundbreaking molecular biology tools [1] and, more recently, to the public domain where the societal and ethical implications and legislation surrounding CRISPR applications are being debated. Some of the potential CRISPR applications that are currently being explored in the laboratory would involve the release of CRISPR genes into confined or open environments—for example, when CRISPR would be used to protect focal bacterial species against phage infections, when it is applied to suppress the spread of antimicrobial resistance or to control vectors of disease [2–4]. One component of the debate surrounding the societal impacts of these applications entails an assessment of the potential risks associated with these strategies (e.g. [5–7]), which requires an understanding of how CRISPR-Cas behaves in an ecological context. In this special issue, we explore this question by examining the evolutionary history of CRISPR-Cas immune systems, where they occur naturally, when they evolve and how this impacts the spread and evolution of other DNA elements. Finally, we return to the question how CRISPR-Cas may be exploited in an ecological context for the benefit of human health, and the ethical challenges that are associated with this.

### (a) CRISPR-Cas adaptive immune systems: a brief overview

CRISPR-Cas adaptive immune systems were discovered around 15 years ago [8–11] and are estimated to exist in approximately 50% of all bacterial genomes and roughly 90% of all archaeal genomes [12]. A CRISPR immunity phenotype is genetically encoded by a so-called CRISPR locus (clustered regularly interspaced short palindromic repeats)—an array of repetitive and unique sequences (repeats and spacers, respectively), both of which are typically around 30 nt in length. Spacers are derived from (foreign) genetic elements, such as plasmids and viruses, and provide immunity to re-infection based on recognition of the cognate sequence (known as 'protospacer') [13–15]. Bacteria or archaea may carry a single linked array of spacers interspersed with repeats (one CRISPR locus) or multiple loci. CRISPR loci can evolve very rapidly owing to insertion of new spacers and the occasional loss of spacers or deletion of the CRISPR locus itself, which can cause very closely related strains to carry unique combinations of spacer sequences, known as a CRISPR allele. The overall length of CRISPR loci will increase and decrease with the acquisition and loss of spacers, and can vary from as little as a single spacer flanked by two repeats to hundreds of spacers and repeats [16]. As new spacers are added at the so-called leader-end of the locus, which is the sequence that contains the CRISPR promoter, CRISPR loci form an inverse chronological record of previous infections from the leader to the trailer end of the locus [17]. The extent

to which different strains share the same spacer sequences in the same order (usually at the trailer end of the CRISPR locus) is commonly used to define related allele groups as a measure of their evolutionary relatedness.

Apart from the genetic CRISPR memory, a functional CRISPR-Cas immune system also requires a set of CRISPR-associated genes (*cas* genes), which encode the protein machinery required for carrying out the immune response [18]. Cas operons vary in their *cas* gene composition and gene synteny, resulting in a classification of CRISPR-Cas systems into two classes, six types and 33 subtypes [19–22]. These diverse CRISPR-Cas variants differ in many of their mechanistic details, which have been discussed elsewhere [23,24], yet also have commonalities in the basic steps of the immune pathway. For example, two Cas proteins—Cas1 and Cas2—are almost invariably part of CRISPR-Cas immune systems and are responsible for inserting new spacer sequences into CRISPR arrays, sometimes assisted by other Cas proteins (reviewed in [15,25]). CRISPR transcripts are processed by either Cas proteins or house-keeping RNases [26], and the resulting processed CRISPR RNAs (crRNAs) are bound by Cas proteins to form a ribonucleoprotein complex that serves to detect and cleave complementary nucleic acid sequences [23].

### (b) Ecology and diversity of CRISPR-Cas immunity

CRISPR-Cas immune systems are unevenly distributed across taxa and environments. For example, only less than half of mesophilic bacteria encode CRISPR-Cas immune systems, compared to over 90% of bacterial thermophiles and archaea (both mesophilic and thermophilic) [11,16,27–29]. Moreover, some uncultured bacterial lineages are virtually devoid of CRISPR-Cas immune systems [30]. A recent computer learning approach suggested that abiotic factors such as oxygen levels and temperature are important predictors of whether microorganisms encode CRISPR-Cas immune systems [31]. However, the ecological drivers of CRISPR distribution remain unclear. In this context, the key question is when CRISPR-Cas is favoured over alternative defence strategies [32]. Experimental, theoretical and correlational studies have suggested a role for viral abundance and diversity [27,28,33,34], direct and indirect fitness costs of CRISPR-Cas immune systems (e.g. autoimmunity, reduced horizontal gene transfer, induced fitness costs, see §1c below) [35–38] and epistasis with other host genes [39,40].

In natural environments, whether that be the human lung, a hot spring or a fermenter, the ecological and evolutionary impact of CRISPR depends on the population-level diversity of immune alleles. The diversity of alleles in the real world may reflect the previous history of interactions and can be used to predict whether infection epidemics will occur in the future. For example, the spread of a virulent virus can eliminate immune diversity from a population by eliminating all susceptible (non-immune) cells, resulting in a selective sweep of an individual strain with a matching CRISPR allele. In the future, this immunodominance across the CRISPR allele might make a population susceptible to the subsequent epidemic spread of a new virus or other mobile element in the population. Starting with a single strain of host or virus, diversity has been shown to evolve in experimental studies [33,41] and to protect bacteria from virus invasion [42]. Data suggest that immune diversity shapes

the evolution of viral pathogens by selecting for recombinant genotypes that are more likely to escape immunity [43], and recombinant microbial CRISPR alleles that are more likely to increase immune profiles of a single strain [44,45].

Interestingly, studies of CRISPR diversity have observed differences across microbial species in different environments. *Leptospirillum* species from acid mine drainage biofilms [43], *Sulfolobus islandicus* strains from a single hot spring [46], Halophilic archaea from a saline pond [47], *Helicobacter cinaedi* [48] and *Pseudomonas aeruginosa* from a single hospital [49] show that a diversity of CRISPR spacers can coexist. By contrast, *P. aeruginosa* within a human lung, or *Prevotella* strains in a single gut sample, are clonal with complete immunodominance at one time and place [49,50]. These differences in diversity among CRISPR populations may result from difference in CRISPR dynamics or demographics (e.g. colonization bottle-necks [51]) and biological constraints (e.g. an inactive spacer acquisition machinery [52]). As CRISPR studies have mainly focused on mechanistic details, our understanding of microbial population dynamics in real populations is surprisingly limited. Additional studies from a diverse range of CRISPR-containing organisms and their viruses, across a range of environments, are needed to identify the basis for the differences in the genetic structure of CRISPR and predict how they impact epidemics and evolutionary dynamics in local populations.

### (c) Coevolution between CRISPR hosts and viruses

Given the detailed molecular understanding of how CRISPR-Cas systems work, one can often predict resistance phenotypes of an individual on the basis of its genotype and that of its phage (or other foreign DNA element). Specifically, while bacteria gain resistance through the acquisition of spacers, phages and other mobile genetic elements can overcome immunity through the acquisition of ‘escape mutations’ in (or sometimes near) the protospacer (e.g. [53,54]). The fascinating details of the biology of CRISPR immunity and phage evasion challenge the classical theoretical framework used to understand host–parasite interactions. In particular, the Lamarckian ability of CRISPR to acquire new spacers allows the bacteria to accumulate a diverse range of resistance alleles against a focal phage [28,55,56]. This multiplicity of resistance within individual bacteria and/or within bacterial populations can overwhelm the evolutionary potential of the phage and drive it to extinction [42,56–58]. Alternative mechanisms of escape are also possible and, recently, a number of different anti-CRISPR (Acr) systems have been described in different phages [59]. These Acr proteins have the ability to downregulate the immunity of CRISPR-Cas systems and allow the phage to exploit CRISPR-resistant bacteria. Interestingly, this immunosuppression has been shown to require the cooperation between multiple phages, where the first phages reduce the efficacy of CRISPR-Cas and allow subsequent infections to exploit the host [60,61].

However, even if phage lack *acr* genes, CRISPR immunity is far from being the silver bullet against phages because it carries several different fitness costs. Some fitness costs are likely to result from the production of an effective interference against invading pathogens and mobile genetic elements [33,62]. Some other costs are associated with self-targeting and auto-immunity [35,36]. Also, when CRISPR targets mobile genetic elements or prophages that

carry some adaptive mutation for the bacteria, CRISPR immunity may also be viewed as a form of self-targeting, because immunity directly harms the bacterial host. For instance, when CRISPR targets beneficial plasmids carrying antibiotic resistance, this immunity can be counter-selected and lead to the loss of CRISPR loci [37,63–65], potentially explaining why many bacteria lack CRISPR-Cas immune systems. Note, however, that apart from a few studies [66], the experimental study of the coevolution between CRISPR immunity and phage evasion is often limited to a few generations. The monitoring of longer coevolution experiments in the laboratory or in natural environments is very welcome to better understand the coevolutionary dynamics driven by CRISPR immunity.

#### (d) Application of CRISPR in an ecological context

While CRISPR-mediated targeting of antibiotics resistance plasmids may be maladaptive for individual bacteria (see above), it was soon realized that this feature of CRISPR systems may be exploited to limit the spread of antibiotics resistance [67–69], and more generally to engineer microbial communities. As the majority of bacteria do not encode end-joining mechanisms to repair double-strand DNA breaks, CRISPR-based technologies are particularly useful for either sequence-specific killing of pathogenic bacteria or removal of accessory genes (e.g. antimicrobial resistance, virulence, etc.). Synthetic CRISPR systems can be delivered to a target bacterium using so-called phagemids, which are replication-deficient bacteriophage particles [67,69], or through conjugative delivery of CRISPR systems [67,70], but research is still in its infancy and considerable challenges are associated with these approaches [3].

CRISPR-mediated ecological engineering has also been explored in higher organisms, where it has raised major interest, given its potential use in pest and invasive species control and reducing vector-borne diseases. Ecological engineering can be achieved by CRISPR-based genome engineering of wild animals followed by introducing these engineered animals into a local wild population. However, this method is likely to be effective only if the target population is relatively small (see [71]). A more powerful, but also riskier approach is to apply synthetic CRISPR-based gene drives [4,72,73]. These elements spread with super-Mendelian inheritance through a population and can disrupt genetic loci or facilitate the spread of genetically linked genes. CRISPR-based gene drives have been successfully used in confined laboratory settings in a range of organisms, including yeast, fruit flies, mosquitoes and, very recently, mice [74–77], and some first insights into the evolutionary dynamics of these drives, including evolution of resistance, are emerging [78–81].

For all these ecological engineering approaches, there is a clear need to better understand the potential risks of these gene drives and whether there is any appetite in society for these types of interventions. For example, what is the durability of these interventions, what are their long-term ecological consequences for (microbial) communities and what countermeasures can be taken to limit or reverse gene drive spread? Being able to predict risks and benefits will be one important factor to get support from local communities, and for ethical and legislative approval. Recognizing the urgent need for new guidelines, scientists from a range of disciplines, including ethicists, social scientists and

biologists, are working together to develop improved safety recommendations for CRISPR-based ecological engineering technologies [6,82,83], and ethical guidelines and the involvement of the public at an early stage in the development of these applications [82,84,85].

## 2. Aims of the theme issue

This theme issue aims to examine when and where CRISPR-Cas immune systems are important, how this impacts the coexistence and coevolution between bacteria and phages and how this ties in with more practical and ethical considerations concerning the application of CRISPR-based technologies in an ecological context. The studies presented in this issue use a wide range of different approaches, from bioinformatics and metagenomics correlational studies to experimental and theoretical analyses, to answer an equally wide range of questions all somehow linked to CRISPR ecology and evolution: from understanding the basic principles of CRISPR evolution in bacteria to community engagement studies to examine if CRISPR-based applications can be applied in the real world to limit the spread of Lyme disease. The theme issue highlights the need to not only understand how CRISPR works in an ecological context, but also to engage with important practical issues when translating this knowledge to real-world applications to improve human health.

## 3. Overview of the papers

This theme issue is broadly divided into four areas that deal, respectively, with (i) the evolutionary history and relative importance of CRISPR-Cas immune systems, (ii) the role of CRISPR-Cas immune systems during bacteria–phage or archaea–virus coevolution in real environments, (iii) understanding the drivers and consequences of coevolution in controlled laboratory environments, and (iv) the way CRISPR may be harnessed to remove infectious diseases from real environments, and the associated considerations that need to be taken into account.

### (a) Evolutionary history and relative importance of CRISPR-Cas immune systems

One of the key questions in the field of CRISPR-Cas is how these sophisticated immune systems evolved in the first place, and how the enormous diversity of CRISPR classes, types and subtypes emerged in this process. Koonin & Makarova [86] apply bioinformatics approaches to bring the evolutionary history of this system in focus. They suggest a key role for a Cas1-related transposase in the early evolution of the ‘Adaptation’ module of the system, followed by evolution of increased complexity through a series of gene duplication and displacement events, recruitment of genes with nuclease activities and in some cases signal transduction genes to form the ‘Interference’ module of the system. While a cloud of uncertainty inevitably surrounds such analyses, their work provides a plausible explanation for the way in which a highly complex adaptive immune system may have evolved. A next obvious question is then, why do not all bacteria and archaea carry this adaptive immune system? This is a recurrent question, and one factor that

clearly matters is how well CRISPR-Cas immune systems perform (in terms of the benefits they provide) relative to alternative immune strategies. For example, many bacteria and archaea encode restriction-modification systems or they can mutate receptors that phages use to attach to the cell surface. So why put up with CRISPR immunity is the question that is raised by Levin and co-workers [87]. Using mathematical modelling, they demonstrate that in theory, the conditions where CRISPR is favoured over these alternative defences are restricted. In other words, microbes have many options available to defend themselves against phages, and CRISPR may simply not always be the best one. However, the benefits of CRISPR are not only determined by the presence or absence of phage, but also by the genetic context of these immune systems, as Bernheim *et al.* [88] show using bioinformatics approaches, and as Chevallereau *et al.* show using experimental manipulations [89]. Specifically, Bernheim *et al.* find evidence for both positive and negative epistatic interactions between CRISPR-Cas subtypes and double-stranded DNA repair pathways in bacteria. This therefore suggests that the distribution of CRISPR-Cas subtypes across bacterial species is shaped by the genetic context, but the underlying mechanism for this association is unclear. Earlier work from the same team demonstrated that one *cas* gene can inhibit the NHEJ pathway, which can be maladaptive, explaining why these systems hardly ever co-occur in the same genome [39]. Chevallereau *et al.* experimentally examined the impact of *mutS*, which is part of the DNA mismatch repair system, on the evolution of CRISPR resistance by *Pseudomonas aeruginosa* against its phage DMS3vir. They found that owing to the resulting increase in mutation supply, bacteria that carried mutations in *mutS* were much more likely to evolve resistance through mutation of the phage receptor, suggesting no selective benefit of carrying CRISPR-Cas immune systems in this host genetic background. Together, these studies show how CRISPR-Cas immune systems may have evolved in the first place, and how natural selection for these systems not only depends on the presence or absence of phage, but also on the presence of alternative defence mechanisms as well as the genetic context of the CRISPR-Cas system.

## (b) Patterns of spacer acquisition in an ecological context

One of the key questions in the field of CRISPR ecology and evolution is what drives the generation and maintenance of CRISPR diversity. Using mathematical modelling, Bradde *et al.* [90] show that spacer acquisition rates can be constrained by a cost of autoimmunity. Their model assumes that higher spacer acquisition rates not only increase the benefits of phage resistance, but also the costs of autoimmunity owing to spacer sampling from the bacterial genome. This results in selection for an optimum spacer acquisition rate that maximizes bacterial survival (a balance of phage resistance and autoimmunity), which in turn depends on factors that influence the infection risk, such as bacterial and phage population sizes. In addition to this theoretical approach, several papers in this issue apply comparative population studies to describe spacer diversity in natural environments, and to identify the factors that drive this diversity. Lopatina *et al.* [91] examine the CRISPR loci of populations of the bacterium *Thermus* in different

geographical locations in Chile, Italy and Russia. They demonstrate that within a single population of the bacterium *Thermus*, more spacers are shared among strains than among distant populations. The authors infer from the pattern that local dynamics between hosts and viruses define the local diversity, although the shared spacers among populations indicate some evidence for gene flow. Similarly, Pauly *et al.* [92] show patterns of persistent local diversity within a single hot spring of *Sulfolobus islandicus*. Pauly also shows that these patterns vary for different viruses, indicating that virus lifestyle impacts immune diversity, and explore different mechanisms of CRISPR escape from different viruses through mutation or virus genome replacement. Finally, Hoikkala *et al.* [93] review the importance of CRISPR-Cas immune systems in aquaculture, and the observed dynamics of coevolution of a bacterial fish pathogen and its phage. They discuss how CRISPR diversity of bacterial strains in these environments can be exploited for tracking the epidemiology of bacterial pathogens, and for phage therapy interventions. Their review highlights how a better understanding of CRISPR diversity may be used to stabilize and shape healthy microbial populations and prevent invasions of bacterial pathogens into aquaculture environments. Collectively, these studies provide novel insights into the generation, maintenance and potential application of spacer diversity in natural environments.

## (c) Coevolution between CRISPR-Cas immune systems and phages in the laboratory

Experiments carried out in controlled environments in the laboratory allow us to explore the details of the coevolutionary processes between CRISPR systems and phages. Common *et al.* [94] carried out a coevolution experiment between *Streptococcus thermophilus* and a lytic phage over 30 days. The monitoring of phenotypic and genotypic changes in both the bacteria and the phage across time confirmed the emergence of an antagonist arms race that often results in the extinction of the phage populations. These extinctions suggest that this coevolution is asymmetric because the phage cannot cope with the accumulation of escape mutations. The existence of a fitness cost associated with most escape mutations was confirmed by a separate study by Chabas *et al.* [95] on the same biological model. In addition, Chabas *et al.* found a dramatic variation in the speed at which the phage can escape different CRISPR-mediated resistance alleles. The resistance induced by some spacers can be easily bypassed by the rapid emergence of escape mutations, while the resistance induced by other spacers is much more durable. The measurement of the phage mutation rate using Luria–Delbruck fluctuation tests allowed Chabas *et al.* to suggest that this variation in resistance durability is likely to be driven by heterogeneity in the mutation rate across different protospacers. The evolution of escape mutations was also studied by Watson *et al.* [96] in two lytic phages infecting CRISPR-resistant *Pectobacterium atrosepticum*. This study contrasted the evolution of the phage against single or multiple spacers. When the resistance was mediated by a single spacer, most escape mutations were owing to a single point mutation in the protospacer adjacent motif (PAM) or the seed sequence. When the resistance is owing to multiple spacers, most phages escaped with deletions in genes encoding structural proteins. These



mutations were viable but affect the morphology and the fitness of the virus, which further support the existence of a cost of escape and an asymmetry in the coevolutionary arms race. The details of phage adaptation were also studied by McKitterick *et al.* [97] in a fascinating system where the phage ICP1 uses a fully functional CRISPR-Cas system to downregulate bacterial immunity of *Vibrio cholerae* mediated by a phage-inducible chromosomal island-like element (PLE). High-throughput sequencing allowed the authors to monitor the acquisition of new spacers against PLE in the CRISPR of ICP1, which provide quantitative resistance against PLE that depends on the number and sequence of spacers. Furthermore, even spacers not targeting the PLE but instead targeting the small chromosome of the bacterium could still reduce bacterial immunity if the chromosomal targeting occurred close to the PLE integration site. Collectively, these different experiments in the laboratory shed a new light on the interference mediated by CRISPR and by the costs associated with different mutations allowing phages to escape bacterial immunity.

#### (d) Implications of CRISPR-Cas for human health and wellbeing

Natural CRISPR-Cas systems can have an important impact on human health by, for example, altering the spread of antimicrobial resistance and virulence genes [98]. However, even though experimental studies show that CRISPR-Cas forms an important barrier for horizontal gene transfer, correlational studies have shown mixed results [63,99]. Here, Shehreen *et al.* [100] examine if the presence of anti-CRISPR genes may impact the correlation between CRISPR-Cas immune systems and antibiotic resistance genes (ARGs). They find a very high degree of variation between bacterial species in the abundance of CRISPR-Cas systems and no correlation with ARG for most species. However, for *P. aeruginosa*, they found a positive correlation between anti-CRISPR and ARG genes, indicating that anti-CRISPRs may facilitate the spread of clinically relevant genes such as those encoding antibiotic resistance or virulence factors in the face of CRISPR-Cas immune systems. These genes may be removed using CRISPR-based ecological engineering technologies. Although this is still a very young field, the developments have moved fast and scientists now have a variety of CRISPR-based tools at hand to alter microbiome composition and function, and use them as next-generation antimicrobials. Ramachandran & Bikard [101] provide a thorough overview of these various CRISPR-based applications to alter the microbiome, including its application in editing bacteria and phage genomes, controlling their gene expression using CRISPRi, killing pathogenic bacteria using CRISPR-based antimicrobials and removing ARGs or other virulence determinants.

While these CRISPR-based ecological engineering approaches have great potential to solve ecological problems, there are many technological, regulatory, societal and ethical challenges. In this issue, de Graeff *et al.* [102] provide a thorough overview of the arguments reported in the scientific literature for and against genome editing in animals, including gene drives, which relate to human health, efficiency, risks and uncertainty, animal welfare, animal dignity, environment and public acceptability. The authors also point out that the ethical debate on genome editing in

animals is predominantly shaped by biomedical and veterinary scientists, and less so by ethicists and social scientists. They argue that involvement of ethicists and social scientists from the very early research stages of technology development may help facilitate responsible governance of animal genome editing applications [103]. Furthermore, there is a need for engagement with the public to address, among others, concerns around equity of access (who will benefit from the new technology?) and the commercialization of the technology (will businesses prioritize profit-making over providing a safe public good?). Buchthal *et al.* [71] provide an excellent example of involving the public in decision-making on CRISPR-based ecological engineering. 'Mice against Ticks' is an exciting new community-guided ecological engineering project aimed at reducing the incidence of tick-borne diseases vectored by mice on the islands of Nantucket and Martha's Vineyard off the US east coast. Their idea is to introduce CRISPR-edited mice that are heritably resistant to ticks and/or tick-borne disease in the islands' local mouse population, which is hypothesized to disrupt the disease transmission cycle. From the stages of conception of the project, the local communities of both islands have been actively involved in all decision-making. Although at a relatively early stage still, successes and challenges from this exciting pilot project will provide future ecological engineering projects with highly valuable insights on how to set up community-driven projects.

## 4. Conclusion and outlook

While our mechanistic understanding of CRISPR-Cas immune systems has raced ahead, our understanding of the evolutionary ecology of these fascinating immune systems is still relatively limited. As outlined above, this issue fills some important gaps in our knowledge of the distribution and impact of CRISPR immune systems, but there are also many outstanding questions that need to be answered. For example, why are there so many different CRISPR-Cas variants, what are their costs and benefits, and how does this depend on the environment? Why do so many bacteria not encode CRISPR-Cas immune systems, and what drives their loss? Why is it that in the laboratory bacteria that encode CRISPR-Cas systems often evolve receptor-based resistance against their phages? Is this because of a lack of ecological realism, widespread anti-CRISPR strategies of phages, or are as yet unknown phage life-history traits important for the evolution of CRISPR resistance? As argued in this introduction, a better understanding of the principles that govern evolution of CRISPR-Cas immune systems and their coevolution with mobile genetic elements can help to inform applications of CRISPR-based technologies in real environments. Apart from this quest for a better understanding of CRISPR ecology and evolution, contributions in this issue also highlight the need for community engagement by CRISPR biologists and participation in debates surrounding the ethics and legislative aspects of these technologies, and provide some excellent examples of how this can be done.

**Data accessibility.** This article has no additional data.

**Competing interests.** We declare we have no competing interests.

**Funding.** E.R.W. is supported by an NERC Independent Research Fellowship (NE/M018350/1) and received funding in the area of CRISPR ecology and evolution from the ERC (ERC-STG-2016-714478—

EVOIMMECH, and EPICUT), BBSRC (BB/N017412/1) and NERC (NE/S001921/1). S.v.H. is supported by a BBSRC Future Leader Fellowship (BB/R010781/1). R.W. is supported for this work by the Cystic Fibrosis Foundation WHITAK16PO, NSF Dimensions of Biodiversity no. 1342876 and an Allen Distinguished Investigator Award.

**Acknowledgements.** The authors thank the Royal Society and all participants of the Hooke meeting on CRISPR ecology and evolution in London, February 2019, and all contributors to the special issue on CRISPR ecology and evolution.

## References

- Adli M. 2018 The CRISPR tool kit for genome editing and beyond. *Nat. Commun.* **9**, 1911. (doi:10.1038/s41467-018-04252-2)
- Selle K, Barrangou R. 2015 CRISPR-based technologies and the future of food science. *J. Food Sci.* **80**, R2367–R2372. (doi:10.1111/1750-3841.13094).
- Pursey E, Sunderhauf D, Gaze WH, Westra ER, van Houte S. 2018 CRISPR–Cas antimicrobials: challenges and future prospects. *PLoS Pathog.* **14**, e1006990. (doi:10.1371/journal.ppat.1006990).
- Noble C, Adlam B, Church GM, Esvelt KM, Nowak MA. 2018 Current CRISPR gene drive systems are likely to be highly invasive in wild populations. *Elife* **7**, e33423. (doi:10.7554/eLife.33423)
- Webber BL, Raghu S, Edwards OR. 2015 Opinion: Is CRISPR-based gene drive a biocontrol silver bullet or global conservation threat? *Proc. Natl Acad. Sci. USA* **112**, 10 565–10 567. (doi:10.1073/pnas.1514258112)
- Akbari OS *et al.* 2015 BIOSAFETY. Safeguarding gene drive experiments in the laboratory. *Science* **349**, 927–929. (doi:10.1126/science.aac7932)
- Brokowski C, Adli M. 2019 CRISPR ethics: moral considerations for applications of a powerful tool. *J. Mol. Biol.* **431**, 88–101. (doi:10.1016/j.jmb.2018.05.044)
- Mojica FJ, Diez-Villasenor C, Garcia-Martinez J, Soria E. 2005 Intervening sequences of regularly spaced prokaryotic repeats derive from foreign genetic elements. *J. Mol. Evol.* **60**, 174–182. (doi:10.1007/s00239-004-0046-3)
- Bolotin A, Quinquis B, Sorokin A, Ehrlich SD. 2005 Clustered regularly interspaced short palindrome repeats (CRISPRs) have spacers of extrachromosomal origin. *Microbiology* **151**, 2551–2561. (doi:10.1099/mic.0.28048-0)
- Pourcel C, Salvignol G, Vergnaud G. 2005 CRISPR elements in *Yersinia pestis* acquire new repeats by preferential uptake of bacteriophage DNA, and provide additional tools for evolutionary studies. *Microbiology* **151**, 653–663. (doi:10.1099/mic.0.27437-0)
- Makarova KS, Grishin NV, Shabalina SA, Wolf YI, Koonin EV. 2006 A putative RNA-interference-based immune system in prokaryotes: computational analysis of the predicted enzymatic machinery, functional analogies with eukaryotic RNAi, and hypothetical mechanisms of action. *Biol. Direct* **1**, 7. (doi:10.1186/1745-6150-1-7)
- Grissa I, Vergnaud G, Pourcel C. 2007 The CRISPRdb database and tools to display CRISPRs and to generate dictionaries of spacers and repeats. *BMC Bioinf.* **8**, 172. (doi:10.1186/1471-2105-8-172)
- Barrangou R, Fremaux C, Deveau H, Richards M, Boyaval P, Moineau S, Romero DA, Horvath P. 2007 CRISPR provides acquired resistance against viruses in prokaryotes. *Science* **315**, 1709–1712. (doi:10.1126/science.1138140)
- Marraffini LA, Sontheimer EJ. 2008 CRISPR interference limits horizontal gene transfer in staphylococci by targeting DNA. *Science* **322**, 1843–1845. (doi:10.1126/science.1165771)
- Amitai G, Sorek R. 2016 CRISPR–Cas adaptation: insights into the mechanism of action. *Nat. Rev. Microbiol.* **14**, 67–76. (doi:10.1038/nrmicro.2015.14)
- Westra ER, Dowling AJ, Broniewski JM, van Houte S. 2016 Evolution and ecology of CRISPR. *Annu. Rev. Ecol. Evol. Syst.* **47**, 307–331. (doi:10.1146/annurev-ecolsys-121415-032428)
- Vale PF, Little TJ. 2010 CRISPR-mediated phage resistance and the ghost of coevolution past. *Proc. R. Soc. B* **277**, 2097–2103. (doi:10.1098/rspb.2010.0055)
- Jansen R, Embden JD, Gaastra W, Schouls LM. 2002 Identification of genes that are associated with DNA repeats in prokaryotes. *Mol. Microbiol.* **43**, 1565–1575. (doi:10.1046/j.1365-2958.2002.02839.x)
- Shmakov SA, Makarova KS, Wolf YI, Severinov KV, Koonin EV. 2018 Systematic prediction of genes functionally linked to CRISPR–Cas systems by gene neighborhood analysis. *Proc. Natl Acad. Sci. USA* **115**, E5307–E5316. (doi:10.1073/pnas.1803440115)
- Shmakov S *et al.* 2017 Diversity and evolution of class 2 CRISPR–Cas systems. *Nat. Rev. Microbiol.* **15**, 169–182. (doi:10.1038/nrmicro.2016.184)
- Koonin EV, Makarova KS, Zhang F. 2017 Diversity, classification and evolution of CRISPR–Cas systems. *Curr. Opin. Microbiol.* **37**, 67–78. (doi:10.1016/j.mib.2017.05.008)
- Makarova KS *et al.* 2015 An updated evolutionary classification of CRISPR–Cas systems. *Nat. Rev. Microbiol.* **13**, 722–736. (doi:10.1038/nrmicro3569)
- van der Oost J, Westra ER, Jackson RN, Wiedenheft B. 2014 Unravelling the structural and mechanistic basis of CRISPR–Cas systems. *Nat. Rev. Microbiol.* **12**, 479–492. (doi:10.1038/nrmicro3279)
- Marraffini LA. 2015 CRISPR–Cas immunity in prokaryotes. *Nature* **526**, 55–61. (doi:10.1038/nature15386)
- Jackson SA, McKenzie RE, Fagerlund RD, Kieper SN, Fineran PC, Brouns SJ. 2017 CRISPR–Cas: adapting to change. *Science* **356**, aa15056. (doi:10.1126/science.aal5056)
- Charpentier E, Richter H, van der Oost J, White MF. 2015 Biogenesis pathways of RNA guides in archaeal and bacterial CRISPR–Cas adaptive immunity. *FEMS Microbiol. Rev.* **39**, 428–441. (doi:10.1093/femsre/fuv023)
- Weinberger AD, Wolf YI, Lobkovsky AE, Gilmore MS, Koonin EV. 2012 Viral diversity threshold for adaptive immunity in prokaryotes. *MBio* **3**, e00456-12. (doi:10.1128/mBio.00456-12)
- Iranzo J, Lobkovsky AE, Wolf YI, Koonin EV. 2013 Evolutionary dynamics of the prokaryotic adaptive immunity system CRISPR–Cas in an explicit ecological context. *J. Bacteriol.* **195**, 3834–3844. (doi:10.1128/JB.00412-13)
- Anderson RE, Brazelton WJ, Baross JA. 2011 Using CRISPRs as a metagenomic tool to identify microbial hosts of a diffuse flow hydrothermal vent viral assemblage. *FEMS Microbiol. Ecol.* **77**, 120–133. (doi:10.1111/j.1574-6941.2011.01090.x)
- Burstein D, Sun CL, Brown CT, Sharon I, Anantharaman K, Probst AJ, Thomas BC, Banfield JF. 2016 Major bacterial lineages are essentially devoid of CRISPR–Cas viral defence systems. *Nat. Commun.* **7**, 10613. (doi:10.1038/ncomms10613)
- Weissman J, Laljani R, Fagan W, Johnson P. 2018 Ecology shapes microbial immune strategy: temperature and oxygen as determinants of the incidence of CRISPR adaptive immunity. *bioRxiv*. (doi:10.1101/326330)
- van Houte S, Buckling A, Westra ER. 2016 Evolutionary ecology of prokaryotic immune mechanisms. *Microbiol. Mol. Biol. Rev.* **80**, 745–763. (doi:10.1128/MMBR.00011-16)
- Westra ER *et al.* 2015 Parasite exposure drives selective evolution of constitutive versus inducible defense. *Curr. Biol.* **25**, 1043–1049. (doi:10.1016/j.cub.2015.01.065)
- Chabas H, van Houte S, Hoyland-Kroghsbo NM, Buckling A, Westra ER. 2016 Immigration of susceptible hosts triggers the evolution of alternative parasite defence strategies. *Proc. R. Soc. B* **283**, 20160721. (doi:10.1098/rspb.2016.0721)
- Veroe RB *et al.* 2013 Cytotoxic chromosomal targeting by CRISPR/Cas systems can reshape bacterial genomes and expel or remodel pathogenicity islands. *PLoS Genet.* **9**, e1003454. (doi:10.1371/journal.pgen.1003454)
- Stern A, Keren L, Wurtzel O, Amitai G, Sorek R. 2010 Self-targeting by CRISPR: gene regulation or autoimmunity? *Trends Genet.* **26**, 335–340. (doi:10.1016/j.tig.2010.05.008)
- Bikard D, Hatoum-Aslan A, Mucida D, Marraffini LA. 2012 CRISPR interference can prevent natural transformation and virulence acquisition during in

- vivo bacterial infection. *Cell Host Microbe* **12**, 177–186. (doi:10.1016/j.chom.2012.06.003)
38. van Sluijs L, van Houte S, van der Oost J, Brouns SJ, Buckling A, Westra ER. In press. Addition systems antagonize bacterial adaptive immunity. *FEMS Microbiol. Lett.*
39. Bernheim A, Calvo-Villamanan A, Basier C, Cui L, Rocha E.PC, Touchon M, Bikard D. 2017 Inhibition of NHEJ repair by type II-A CRISPR–Cas systems in bacteria. *Nat. Commun.* **8**, 2094. (doi:10.1038/s41467-017-02350-1)
40. Levy A, Goren MG, Yosef I, Auster O, Manor M, Amitai G, Edgar R, Qimron U, Sorek R. 2015 CRISPR adaptation biases explain preference for acquisition of foreign DNA. *Nature* **520**, 505–510. (doi:10.1038/nature14302)
41. Paez-Espino D, Sharon I, Morovic W, Stahl B, Thomas BC, Barrangou R, Banfield JF. 2015 CRISPR immunity drives rapid phage genome evolution in *Streptococcus thermophilus*. *MBio* **6**, e00262–15. (doi:10.1128/mBio.00262-15)
42. van Houte S *et al.* 2016 The diversity-generating benefits of a prokaryotic adaptive immune system. *Nature* **532**, 385–388. (doi:10.1038/nature17436)
43. Andersson AF, Banfield JF. 2008 Virus population dynamics and acquired virus resistance in natural microbial communities. *Science* **320**, 1047–1050. (doi:10.1126/science.1157358)
44. Held NL, Herrera A, Whitaker RJ. 2013 Reassortment of CRISPR repeat-spacer loci in *Sulfolobus islandicus*. *Environ. Microbiol.* **15**, 3065–3076. (doi:10.1111/1462-2920.12146)
45. Tyson GW, Banfield JF. 2008 Rapidly evolving CRISPRs implicated in acquired resistance of microorganisms to viruses. *Environ. Microbiol.* **10**, 200–207. (doi:10.1111/j.1462-2920.2007.01444.x)
46. Held NL, Herrera A, Cadillo-Quiroz H, Whitaker RJ. 2010 CRISPR associated diversity within a population of *Sulfolobus islandicus*. *PLoS ONE* **5**, e0012988. (doi:10.1371/journal.pone.0012988)
47. Emerson JB, Andrade K, Thomas BC, Norman A, Allen EE, Heidelberg KB, Banfield JF. 2013 Virus-host and CRISPR dynamics in Archaea-dominated hypersaline Lake Tyrrell, Victoria, Australia. *Archaea* **2013**, 370871. (doi:10.1155/2013/370871)
48. Tomida J, Morita Y, Shibayama K, Kikuchi K, Sawa T, Akaike T, Kawamura Y. 2017 Diversity and microevolution of CRISPR loci in *Helicobacter cinaedi*. *PLoS ONE* **12**, e0186241. (doi:10.1371/journal.pone.0186241)
49. England WE, Kim T, Whitaker RJ. 2018 Metapopulation structure of CRISPR–Cas immunity in *Pseudomonas aeruginosa* and its viruses. *mSystems* **3**, e00075–18. (doi:10.1128/mSystems.00075-18)
50. Devoto AE *et al.* In press. Megaphages infect *Prevotella* and variants are widespread in gut microbiomes. *Nat. Microbiol.* (doi:10.1038/s41564-018-0338-9)
51. Common J, Westra ER. 2019 CRISPR evolution and bacteriophage persistence in the context of population bottlenecks. *RNA Biol.* **5**, 1–7. (doi:10.1080/15476286.2019.1578608)
52. Touchon M, Rocha EP. 2010 The small, slow and specialized CRISPR and anti-CRISPR of *Escherichia* and *Salmonella*. *PLoS ONE* **5**, e11126. (doi:10.1371/journal.pone.0011126)
53. Semenova E, Jore MM, Datsenko KA, Semenova A, Westra ER, Wanner B, van der Oost J, Brouns SJ, Severinov K. 2011 Interference by clustered regularly interspaced short palindromic repeat (CRISPR) RNA is governed by a seed sequence. *Proc. Natl Acad. Sci. USA* **108**, 10 098–10 103. (doi:10.1073/pnas.1104144108)
54. Deveau H, Barrangou R, Garneau JE, Labonte J, Fremaux C, Boyaval P, Romero DA, Horvath P, Moineau S. 2008 Phage response to CRISPR-encoded resistance in *Streptococcus thermophilus*. *J. Bacteriol.* **190**, 1390–1400. (doi:10.1128/JB.01412-07)
55. Levin BR, Moineau S, Bushman M, Barrangou R. 2013 The population and evolutionary dynamics of phage and bacteria with CRISPR-mediated immunity. *PLoS Genet.* **9**, e1003312. (doi:10.1371/journal.pgen.1003312)
56. Childs LM, England WE, Young MJ, Weitz JS, Whitaker RJ. 2014 CRISPR-induced distributed immunity in microbial populations. *PLoS ONE* **9**, e101710. (doi:10.1371/journal.pone.0101710)
57. Chabas H, Lion S, Nicot A, Meaden S, van Houte S, Moineau S, Wahl LM, Westra ER, Gandon S. 2018 Evolutionary emergence of infectious diseases in heterogeneous host populations. *PLoS Biol.* **16**, e2006738. (doi:10.1371/journal.pbio.2006738)
58. Westra ER, Sunderhauf D, Landsberger M, Buckling A. 2017 Mechanisms and consequences of diversity-generating immune strategies. *Nat. Rev. Immunol.* **17**, 719–728. (doi:10.1038/nri.2017.78)
59. Bondy-Denomy J, Pawluk A, Maxwell KL, Davidson AR. 2013 Bacteriophage genes that inactivate the CRISPR/Cas bacterial immune system. *Nature* **493**, 429–432. (doi:10.1038/nature11723)
60. Borges AL, Zhang JY, Rollins MF, Osuna BA, Wiedenheft B, Bondy-Denomy J. 2018 Bacteriophage cooperation suppresses CRISPR–Cas3 and Cas9 immunity. *Cell* **174**, 917–925. (doi:10.1016/j.cell.2018.06.013)
61. Landsberger M, Gandon S, Meaden S, Rollie C, Chevallereau A, Chabas H, Buckling A, Westra ER, van Houte S. 2018 Anti-CRISPR phages cooperate to overcome CRISPR–Cas immunity. *Cell* **174**, 908–916. (doi:10.1016/j.cell.2018.05.058)
62. Vale PF, Lafforgue G, Gatchitch F, Gardan R, Moineau S, Gandon S. 2015 Costs of CRISPR–Cas-mediated resistance in *Streptococcus thermophilus*. *Proc. R. Soc. B* **282**, 20151270. (doi:10.1098/rspb.2015.1270)
63. Palmer KL, Gilmore MS. 2010 Multidrug-resistant enterococci lack CRISPR–cas. *MBio* **1**, e00227–10. (doi:10.1128/mBio.00227-10)
64. Jiang W, Maniv I, Arain F, Wang Y, Levin BR, Marraffini LA. 2013 Dealing with the evolutionary downside of CRISPR immunity: bacteria and beneficial plasmids. *PLoS Genet.* **9**, e1003844. (doi:10.1371/journal.pgen.1003844)
65. Gandon S, Vale PF. 2014 The evolution of resistance against good and bad infections. *J. Evol. Biol.* **27**, 303–312. (doi:10.1111/jeb.12291)
66. Paez-Espino D, Morovic W, Sun CL, Thomas BC, Ueda K, Stahl B, Barrangou R, Banfield JF. 2013 Strong bias in the bacterial CRISPR elements that confer immunity to phage. *Nat. Commun.* **4**, 1430. (doi:10.1038/ncomms2440)
67. Citorik RJ, Mimee M, Lu TK. 2014 Sequence-specific antimicrobials using efficiently delivered RNA-guided nucleases. *Nat. Biotechnol.* **32**, 1141–1145. (doi:10.1038/nbt.3011)
68. Goma AA, Klumpe HE, Luo ML, Selle K, Barrangou R, Beisel CL. 2013 Programmable removal of bacterial strains by use of genome-targeting CRISPR–Cas systems. *MBio* **5**, e00928–13. (doi:10.1128/mBio.00928-13)
69. Bikard D, Euler CW, Jiang W, Nussenzweig PM, Goldberg GW, Duportet X, Fischetti VA, Marraffini LA. 2014 Exploiting CRISPR–Cas nucleases to produce sequence-specific antimicrobials. *Nat. Biotechnol.* **32**, 1146–1150. (doi:10.1038/nbt.3043)
70. Dong H, Xiang H, Mu D, Wang D, Wang T. 2019 Exploiting a conjugative CRISPR/Cas9 system to eliminate plasmid harbouring the *mcr-1* gene from *Escherichia coli*. *Int. J. Antimicrob. Agents* **53**, 1–8. (doi:10.1016/j.ijantimicag.2018.09.017)
71. Buchthal J, Evans SW, Lunshof J, Telford III SR, Esvelt KM. 2019 Mice Against Ticks: an experimental community-guided effort to prevent tick-borne disease by altering the shared environment. *Phil. Trans. R. Soc. B* **374**, 20180105. (doi:10.1098/rstb.2018.0105)
72. Esvelt KM, Smidler AL, Catteruccia F, Church GM. 2014 Concerning RNA-guided gene drives for the alteration of wild populations. *Elife* **3**, e03401. (doi:10.7554/eLife.03401)
73. Godfray H.CJ, North A, Burt A. 2017 How driving endonuclease genes can be used to combat pests and disease vectors. *BMC Biol.* **15**, 81. (doi:10.1186/s12915-017-0420-4)
74. DiCarlo JE, Chavez A, Dietz SL, Esvelt KM, Church GM. 2015 Safeguarding CRISPR–Cas9 gene drives in yeast. *Nat. Biotechnol.* **33**, 1250–1255. (doi:10.1038/nbt.3412)
75. Gantz VM, Jasinskiene N, Tatarenkova O, Fazekas A, Macias VM, Bier E, James AA. 2015 Highly efficient Cas9-mediated gene drive for population modification of the malaria vector mosquito *Anopheles stephensi*. *Proc. Natl Acad. Sci. USA* **112**, E6736–E6743. (doi:10.1073/pnas.1521077112)
76. Gantz VM, Bier E. 2015 Genome editing. The mutagenic chain reaction: a method for converting heterozygous to homozygous mutations. *Science* **348**, 442–444. (doi:10.1126/science.aaa5945)
77. Grunwald HA, Gantz VM, Poplawski G, Xu XS, Bier E, Cooper KL. 2019 Super-Mendelian inheritance mediated by CRISPR–Cas9 in the female mouse germline. *Nature* **566**, 105–109. (doi:10.1038/s41586-019-0875-2)
78. Champer J, Liu J, Oh SY, Reeves R, Luthra A, Oakes N, Clark AG, Messer PW. 2018 Reducing resistance allele



- formation in CRISPR gene drive. *Proc. Natl Acad. Sci. USA* **115**, 5522–5527. (doi:10.1073/pnas.1720354115)
79. Unckless RL, Clark AG, Messer PW. 2017 Evolution of resistance against CRISPR/Cas9 gene drive. *Genetics* **205**, 827–841. (doi:10.1534/genetics.116.197285)
  80. Noble C, Olejarz J, Esvelt K, Church G, Nowak M. 2017 Evolutionary dynamics of CRISPR gene drives. *Sci. Adv.* **3**, e1601964. (doi:10.1126/sciadv.1601964)
  81. Kyrou K, Hammond AM, Galizi R, Kranjc N, Burt A, Beaghton AK, Nolan T, Crisanti A. 2018 A CRISPR–Cas9 gene drive targeting *doublesex* causes complete population suppression in caged *Anopheles gambiae* mosquitoes. *Nat. Biotechnol.* **36**, 1062–1066. (doi:10.1038/nbt.4245)
  82. James S *et al.* 2018 Pathway to deployment of gene drive mosquitoes as a potential biocontrol tool for elimination of malaria in sub-Saharan Africa: recommendations of a Scientific Working Group. *Am. J. Trop. Med. Hyg.* **98**(Suppl. 6), 1–49. (doi:10.4269/ajtmh.18-0083)
  83. Min J, Smidler AL, Najjar D, Esvelt KM. 2018 Harnessing gene drive. *J. Responsible Innov.* **5**(Suppl. 1), S40–S65. (doi:10.1080/23299460.2017.1415586)
  84. National Academies of Sciences Engineering and Medicine. 2016 *Gene drives on the horizon: advancing science, navigating uncertainty, and aligning research with public values*. Washington, DC: The National Academies Press.
  85. Benedict M *et al.* 2014 Guidance framework for testing of genetically modified mosquitoes. FNIH, WHO, TDR. See <https://manualzz.com/doc/18478798/guidance-framework-for-testing-of-genetically-modified-mosquitoes>.
  86. Koonin EV, Makarova KS. 2019 Origins and evolution of CRISPR–Cas systems. *Phil. Trans. R. Soc. B* **374**, 20180087. (doi:10.1098/rstb.2018.0087)
  87. Gurney J, Pleška M, Levin BR. 2019 Why put up with immunity when there is resistance: an excursion into the population and evolutionary dynamics of restriction–modification and CRISPR–Cas. *Phil. Trans. R. Soc. B* **374**, 20180096. (doi:10.1098/rstb.2018.0096)
  88. Bernheim A, Bikard D, Touchon M, Rocha EPC. 2019 A matter of background: DNA repair pathways as a possible cause for the sparse distribution of CRISPR–Cas systems in bacteria. *Phil. Trans. R. Soc. B* **374**, 20180088. (doi:10.1098/rstb.2018.0088)
  89. Chevallereau A, Meaden S, van Houte S, Westra ER, Rollic C. 2019 The effect of bacterial mutation rate on the evolution of CRISPR–Cas adaptive immunity. *Phil. Trans. R. Soc. B* **374**, 20180094. (doi:10.1098/rstb.2018.0094)
  90. Bradde S, Mora T, Walczak AM. 2019 Cost and benefits of clustered regularly interspaced short palindromic repeats spacer acquisition. *Phil. Trans. R. Soc. B* **374**, 20180095. (doi:10.1098/rstb.2018.0095)
  91. Lopatina A, Medvedeva S, Artamonova D, Kolesnik M, Sitnik V, Ispolatov Y, Severinov K. 2019 Natural diversity of CRISPR spacers of *Thermus*: evidence of local spacer acquisition and global spacer exchange. *Phil. Trans. R. Soc. B* **374**, 20180092. (doi:10.1098/rstb.2018.0092)
  92. Pauly MD, Bautista MA, Black JA, Whitaker RJ. 2019 Diversified local CRISPR–Cas immunity to viruses of *Sulfolobus islandicus*. *Phil. Trans. R. Soc. B* **374**, 20180093. (doi:10.1098/rstb.2018.0093)
  93. Hoikkala V, Almeida GMF, Laanto E, Sundberg L-R. 2019 Aquaculture as a source of empirical evidence for coevolution between CRISPR–Cas and phage. *Phil. Trans. R. Soc. B* **374**, 20180100. (doi:10.1098/rstb.2018.0100)
  94. Common J, Morley D, Westra ER, van Houte S. 2019 CRISPR–Cas immunity leads to a coevolutionary arms race between *Streptococcus thermophilus* and lytic phage. *Phil. Trans. R. Soc. B* **374**, 20180098. (doi:10.1098/rstb.2018.0098)
  95. Chabas H, Nicot A, Meaden S, Westra ER, Tremblay DM, Pradier L, Lion S, Moineau S, Gandon S. 2019 Variability in the durability of CRISPR–Cas immunity. *Phil. Trans. R. Soc. B* **374**, 20180097. (doi:10.1098/rstb.2018.0097)
  96. Watson BNJ, Easingwood RA, Tong B, Wolf M, Salmond GPC, Staals RHJ, Bostina M, Fineran PC. 2019 Different genetic and morphological outcomes for phages targeted by single or multiple CRISPR–Cas spacers. *Phil. Trans. R. Soc. B* **374**, 20180090. (doi:10.1098/rstb.2018.0090)
  97. McKitterick AC, LeGault KN, Angermeyer A, Alam M, Seed KD. 2019 Competition between mobile genetic elements drives optimization of a phage-encoded CRISPR–Cas system: insights from a natural arms race. *Phil. Trans. R. Soc. B* **374**, 20180089. (doi:10.1098/rstb.2018.0089)
  98. Hatoum-Aslan A, Marraffini LA. 2014 Impact of CRISPR immunity on the emergence and virulence of bacterial pathogens. *Curr. Opin. Microbiol.* **17C**, 82–90. (doi:10.1016/j.mib.2013.12.001)
  99. Gophna U, Kristensen DM, Wolf YI, Popa O, Drevet C, Koonin EV. 2015 No evidence of inhibition of horizontal gene transfer by CRISPR–Cas on evolutionary timescales. *ISME J.* **9**, 2021–2027. (doi:10.1038/ismej.2015.20)
  100. Shehreen S, Chyou T-y, Fineran PC, Brown CM. 2019 Genome-wide correlation analysis suggests different roles of CRISPR–Cas systems in the acquisition of antibiotic resistance genes in diverse species. *Phil. Trans. R. Soc. B* **374**, 20180384. (doi:10.1098/rstb.2018.0384)
  101. Ramachandran G, Bikard D. 2019 Editing the microbiome the CRISPR way. *Phil. Trans. R. Soc. B* **374**, 20180103. (doi:10.1098/rstb.2018.0103)
  102. de Graeff N, Jongsma KR, Johnston J, Hartley S, Bredenoord AL. 2019 The ethics of genome editing in non-human animals: a systematic review of reasons reported in the academic literature. *Phil. Trans. R. Soc. B* **374**, 20180106. (doi:10.1098/rstb.2018.0106)
  103. Hartley S, Gillund F, van Hove L, Wickson F. 2016 Essential features of responsible governance of agricultural biotechnology. *PLoS Biol.* **14**, e1002453. (doi:10.1371/journal.pbio.1002453)