



Research

Cite this article: Knoch S, Whiteside MA, Madden JR, Rose PE, Fawcett TW. 2022 Hot-headed peckers: thermographic changes during aggression among juvenile pheasants (*Phasianus colchicus*). *Phil. Trans. R. Soc. B* **377**: 20200442.
<https://doi.org/10.1098/rstb.2020.0442>

Received: 23 June 2021
Accepted: 28 October 2021

One contribution of 19 to a theme issue ‘The centennial of the pecking order: current state and future prospects for the study of dominance hierarchies’.

Subject Areas:
behaviour, cognition, physiology

Keywords:
aggression, dominance hierarchy, infrared thermography, pecking order, social defeat, stress-induced hyperthermia

Author for correspondence:
Tim W. Fawcett
e-mail: t.w.fawcett@exeter.ac.uk

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5742174>.

Hot-headed peckers: thermographic changes during aggression among juvenile pheasants (*Phasianus colchicus*)

Sophia Knoch^{1,2}, Mark A. Whiteside^{1,3}, Joah R. Madden¹, Paul E. Rose¹ and Tim W. Fawcett¹

¹Centre for Research in Animal Behaviour (CRAB), Washington Singer Laboratories, University of Exeter, Exeter EX4 4QG, UK

²Institute of Psychology, University of Freiburg, Engelbergerstr. 41, 79085 Freiburg, Germany

³School of Biological and Marine Sciences, University of Plymouth, Drake Circus, Plymouth PL4 8AA, UK

MAW, 0000-0001-7212-7279; JRM, 0000-0002-0691-0967; PER, 0000-0002-5375-8267; TWF, 0000-0001-6337-901X

In group-living vertebrates, dominance status often covaries with physiological measurements (e.g. glucocorticoid levels), but it is unclear how dominance is linked to dynamic changes in physiological state over a shorter, behavioural timescale. In this observational study, we recorded spontaneous aggression among captive juvenile pheasants (*Phasianus colchicus*) alongside infrared thermographic measurements of their external temperature, a non-invasive technique previously used to examine stress responses in non-social contexts, where peripheral blood is redirected towards the body core. We found low but highly significant repeatability in maximum head temperature, suggesting individually consistent thermal profiles, and some indication of lower head temperatures in more active behavioural states (e.g. walking compared to resting). These individual differences were partly associated with sex, females being cooler on average than males, but unrelated to body size. During pairwise aggressive encounters, we observed a non-monotonic temperature change, with head temperature dropping rapidly immediately prior to an attack and increasing rapidly afterwards, before returning to baseline levels. This nonlinear pattern was similar for birds in aggressor and recipient roles, but aggressors were slightly hotter on average. Our findings show that aggressive interactions induce rapid temperature changes in dominants and subordinates alike, and highlight infrared thermography as a promising tool for investigating the physiological basis of pecking orders in galliforms.

This article is part of the theme issue ‘The centennial of the pecking order: current state and future prospects for the study of dominance hierarchies’.

1. Introduction

Group-living animals often form dominance hierarchies in which some individuals consistently outrank others [1], gaining preferential access to food, mates or other resources [2]. Since Schjelderup-Ebbe’s [3] pioneering observations on ‘pecking orders’ in domestic chickens (*Gallus gallus domesticus*), there has been continued interest in how these hierarchies are formed and maintained [4,5]. Researchers have studied the role of pre-existing behavioural and physiological differences between individuals [6,7], the reinforcement of previous contest outcomes [8] through phenomena such as winner and loser effects [9–11] and bystander effects [12], and the impacts of dominance interactions on physiology and cognition [13]. In social vertebrates, dominance status has been linked to differences in circulating glucocorticoid levels [14] and even symptoms of chronic psychosocial stress [15]. However, the precise relationship between stress physiology and dominance status is variable and context-dependent [15–22]. Due to the complex, reciprocal links between hormones

and behaviour, it is often unclear to what extent physiological differences determine dominance and *vice versa* [23].

One key issue is how organisms respond to their experiences in individual dominance encounters [10]. Assays of blood and brain tissue after staged dyadic contests in the laboratory have shown that winning and losing experiences may be associated with neuroendocrine changes in the hypothalamic–pituitary–adrenocortical (HPA) system, involving serotonin and glucocorticoids [24]. In rainbow trout (*Oncorhynchus mykiss*) and green anoles (*Anolis carolinensis*), for example, aggressive interactions between males lead to elevated glucocorticoids in both winners and losers, but this increase tends to be much longer lasting for losers [25,26]. Experimental administration of exogenous glucocorticoids appears to have differing effects depending on the length of exposure: acute doses tend to promote aggression, while prolonged treatment may reduce it [23,24].

Notwithstanding the value of these approaches, blood and tissue sampling has some ethical and practical limitations, because it is invasive and requires capture and handling (or even sacrifice) of the animal. This creates a time lag from the event of interest to the sampling point [27] and acts as an additional stressor, directly affecting the physiological measures of interest [14,28,29] and potentially causing injury and infections. Time lags are also a problem for faecal sampling, a non-invasive method commonly used in field studies [29]. While useful for examining longer term associations between stress physiology and dominance rank [14,15,30], the timing of defecation by the animal is beyond the experimenter's control and faecal glucocorticoids reflect stress levels several hours earlier [27,29], making it difficult to link them to specific events experienced by the animal. Assessing the impact of individual dominance encounters, which are typically brief and often happen in quick succession, requires measurements of physiological change that are not subject to time lags, and ideally are non-invasive.

Changes in body temperature offer another potential route to examining immediate physiological responses to stressors [27]. Stress-induced hyperthermia (SIH), also known as 'psychogenic fever', is a common response in endothermic animals to perceived threats in which sympathetically mediated vasoconstriction redirects peripheral blood flow towards the body core, raising core temperature [31,32]. Humans, for instance, show SIH prior to exams [33,34]. In research on non-human animals, where body temperature can be monitored using implanted devices [35], SIH has been found to correlate with a faster heart rate [36] and increased glucocorticoid levels [24] in response to a stressor. For instance, core body temperature, heart rate and plasma corticosterone increased in laboratory mice (*Mus musculus*) when handled or injected [37–39], with some indication of a stronger response in strains selectively bred for high aggression [38].

SIH can also be triggered by social stressors. Several studies have shown that social defeat leads to increased core body temperature in laboratory rodents [40–42]. In captive male great tits (*Parus major*), core body temperature (measured using a probe inserted down the throat) remained elevated for 1 day after experiencing a social defeat, and defeated males also tended to avoid social interaction with conspecifics during this period [43]. These studies induced social defeat using a resident–intruder paradigm, whereby an 'intruding' individual is introduced to the home cage of a 'resident' individual, with the latter almost always winning [44]. Such studies have focused on dyadic

interactions under enforced conditions. We know far less about physiological responses to the more natural dominance interactions that occur spontaneously in larger groups. Implanted devices have been used to detect heart rate changes during aggression in free-roaming greylag geese (*Anser anser*) [45], but, as with studies of core body temperature, the practical and ethical challenges associated with invasive procedures limit their applicability [46].

While elevating core body temperature, SIH simultaneously results in a rapid drop (within seconds) in surface body temperature [47,48]. In animals with exposed skin areas, such as the periophthalmic ring in birds, this temperature drop can be detected through remote thermal imaging using infrared cameras [49], as demonstrated in a series of studies on poultry [32,50,51], wild passerines [52–54] and gamebirds [55]. A key advantage of infrared thermography (IRT) is that it avoids the use of implants or other invasive probes needed to measure core body temperature, which require capture, handling and other procedures that themselves are likely to induce a stress response [56,57]. In primates, IRT has revealed rapid physiological responses to social stressors: nasal skin temperature in captive rhesus macaques (*Macaca mulatta*) dropped within seconds of viewing a video clip of an aggressively raging conspecific [58], while wild chimpanzees (*Pan troglodytes*) showed changes in both nose temperature and ear temperature after hearing aggressive vocalizations from a rival group [59]. In birds, studies using IRT to measure stress responses have focused primarily on non-social stressors such as disease [60], heat [35], capture [32] and food removal [51]. General effects of group-level housing conditions (e.g. stocking density) have also been found [61], but there have been no studies to our knowledge investigating responses to direct interactions with conspecifics.

To address this gap, we recorded acts of aggressive dominance behaviour occurring naturally in groups of captive-reared juvenile pheasants (*Phasianus colchicus*), alongside detailed infrared thermographic measurements of their head temperature. Like other galliforms, juvenile pheasants have areas of naturally bare skin around the eye and ear with a high density of blood vessels [62], making them highly suitable for IRT studies. Pheasants are precocial, so in captivity a large number of chicks can be hatched on the same day and reared under standardized conditions without their parents [63], thereby eliminating differences in age and parental care. In the wild, pheasants exhibit harem defence polygyny, with dominant males maintaining control of territories (and access to females) over a prolonged period [63]. An individual adult male's social rank strongly influences his mating success [64]. In captivity, pheasant chicks are aggressive towards one another and sexual segregation emerges within the first few weeks of life, perhaps driven by female avoidance of aggressive males [65].

Our main aim was to examine whether IRT reveals dynamic physiological changes occurring during dominance encounters. We tested three key hypotheses. First, in line with evidence across taxa that stress physiology is moderately repeatable within individuals [66], we predicted that head temperatures in pheasants would show consistent individual differences, potentially linked to other measurable characteristics such as sex and size. Alternatively, within-individual changes in temperature associated with different behavioural states (as seen in chickens [32]) might overwhelm between-individual differences, leading to low repeatability. Second, based on the

suggestion that stress physiology covaries with behavioural differences such as proactive/reactive ‘coping styles’ [67], we hypothesized that individual differences in head temperature would predict the roles that individuals adopt in dominance encounters with their group mates, in terms of whether they are the aggressor or the recipient of aggression. Third, we predicted that during an aggressive encounter pheasants would show a drop in surface head temperature, reflecting a stress-induced hyperthermic response in which blood is redirected from the periphery to the body core [31]. We expected that SIH would occur in both aggressors and recipients, but that the response in recipients would be more pronounced and/or longer lasting (as in [25,26,68]).

2. Methods

(a) Subjects and housing

We used 126 juvenile pheasants (*Phasianus colchicus*), aged 6–7 weeks at the time of our study, that were being reared at the Rothamsted Research farm at North Wyke, Devon, UK. The birds were a mix of full-sibs, half-sibs and unrelated individuals that had hatched in artificial incubators from eggs collected from pens of freely mating polygynandrous adults. On 24 May 2018, when they were 1-day old, the chicks had been randomly allocated to four mixed-sex groups (three $n = 32$, one $n = 30$) and housed in indoor pens (1 × 2 m floor area) that contained a heat lamp, feeder, drinker and assorted branches. From 3 weeks old, each group could also access an outdoor run (12 × 4 m) with additional food, water and perching locations. Age- and nutrient-specific food (Keeper’s Choice, Norfolk, UK) and water were available *ad libitum* throughout rearing. Each pheasant bore a uniquely numbered patagial tag for individual identification. At 9 weeks old (i.e. after our study), the birds were sexed (via plumage traits), weighed (using a spring balance; precision = 5 g) and measured (tarsus length and wing length, using callipers; precision = 0.1 mm) before being released into an open-topped woodland pen covering approximately 4000 m², as part of other studies into their ecology and cognition. Mass at release was highly correlated with mass measured at 6 weeks ($r = 0.90$, $t_{124} = 23.1$, $p < 0.001$), close to the point of our study.

(b) Procedure

On 5 and 13 July 2018, when they were 6 to 7 weeks old, the pheasants in each group were confined to their indoor pen (after removal of the feeder) for up to 30 min as part of routine husbandry procedures. Studies in other galliforms have shown that physiological stress responses are readily detectable by this age [69,70]. We filmed each group once during this confinement period, using two cameras fixed to a tripod positioned approximately 1 m outside the pen: a FLIR T530 thermal video camera to record the birds’ external temperature, plus a Sony HDR CX625 video camera to identify individuals from their numbered patagial tags (which were not visible in the infrared thermal footage). The position of the tripod ensured that all thermal measurements were taken from a distance of 1.0–2.3 m. Recordings from the two cameras were synchronized via hand waving to signal the start of the observation period, which lasted for 13 min for each group.

(c) Video coding

All four videos were coded by one member of the research team (SK), who followed each bird over the 13 min of footage to obtain two sets of behavioural observations. First, for the *baseline* observations, every 30 s we recorded the bird’s behavioural state,

Table 1. Ethogram used to classify behavioural states of captive juvenile pheasants at 30 s time intervals, adapted from [71,72].

behaviour	description
resting	Body flush with substrate, wings tucked and head either upright or relaxed. Eyes open or closed.
standing	Balanced upright on both feet with legs extended. No body movement but head may be moving. Head upright or relaxed and eyes open or closed.
walking	Making more than one step with feet in one direction. Head upright.
exploratory pecking	Using beak to peck other pheasants gently, without aggression.
aggression	Threatening, pecking or fighting with other pheasants (table 2).
preening	Using beak to clean wings and feathers. Also includes feather ruffling and wing stretching.
foraging	Head lowered, with beak pecking or scratching at the floor.
other	Other behaviours not covered by any of the above descriptions.

using the ethogram in table 1. Second, for the *aggression* observations, we identified any aggressive encounters with another bird. For each encounter, we noted the identities of both birds involved, the roles they played (aggressor or recipient), the outcome (winner or loser) and the type of aggression (threat, peck or fight; table 2). The bird that initiated the aggressive encounter was classified as the ‘aggressor’, while the bird to which the aggressive behaviour was directed was the ‘recipient’. The bird that performed the last aggressive act in the encounter was designated the ‘winner’, while if a bird retreated from its opponent and did not retaliate it was designated the ‘loser’. In all encounters we observed, the aggressor ‘won’ and the recipient ‘lost’ (i.e. did not retaliate), so aggressor/recipient roles can be considered synonymous with win/lose outcomes in our analysis.

(d) Thermal measurements

At the same time points as the behavioural observations described above, we used the box selection tool in *FLIR Tools* software version 5.13 (FLIR Systems, Inc. 2015) to pinpoint the maximum temperature on the pheasant’s head (figure 1).

(i) Baseline measurements

Baseline measurements were taken at 30 s intervals throughout the 13 min video, except when the bird was hidden from view or could not be reliably identified, or its head was not at a lateral angle to the camera. We also excluded any measurements taken within 20 s of an aggressive encounter involving that bird. At the same 30 s intervals, we took spot measurements of a suspended white plastic drinker filled with water (figure 1), which provided a stable background reference temperature.




Figure 1. Infrared thermographic image taken from *FLIR Tools*, showing an aggressive encounter between two juvenile pheasants (middle right of the image). The scale at the top indicates the colour coding of temperatures, while the red, upwards-pointing triangles automatically pinpoint the maximum temperature within a selection box drawn manually around each pheasant's head. Here, the aggressor on the left (Bx1), with a maximum head temperature of 37.5°C, has just delivered an aggressive peck to the recipient on the right (Bx2), who has a maximum head temperature of 35.7°C. The background reference temperature is 26.5°C, taken from a white plastic drinker filled with water (Sp1), suspended above the pen floor (top left of image). (Online version in colour.)

Table 2. Ethogram used to classify the types of aggressive encounter between captive juvenile pheasants, in order of increasing intensity. Adapted from [73].

behaviour	description
threat	Bird raises head and neck rapidly, stares at opponent and appears ready to deliver an aggressive peck. Usually face to face with opponent.
peck	Rapid downward stabbing motion with the beak, directed towards the head of another bird.
fight	Birds stand directly in front of each other, necks and heads raised at the same level, and deliver vigorous kicks to the opponent.

(ii) Aggression measurements

During aggressive encounters, we recorded the maximum head temperature of both birds every second from 5 s before to 20 s after the aggressive behaviour occurred, henceforth referred to as the moment of 'attack' (even if this involved no physical contact, as in the case of threats). This period was chosen because previous literature suggests that thermal values drop within 10–20 s after exposure to a stressor [54,58]. To reduce data overlap when aggressive encounters occurred in bursts, we only included encounters for which a minimum of 20 s had passed since that bird's last encounter.

(e) Statistical analysis

All data processing and analysis were conducted in *R* v. 3.6.3 [74]. Body condition at release was calculated as mass (g) divided by tarsus length (mm) cubed. To analyse variation in the baseline thermal measurements, we fitted a linear mixed-effects model (LMM) in package *lme4* [75], with a random effect that allowed varying intercepts across individual pheasants to account for the non-independence of repeated measurements from the same individual; group was modelled as a fixed effect rather than a random effect

because there were only four replicates. We used the package *rptR* [76] to calculate the repeatability (intra-class correlation coefficient) of maximum head temperature within individuals and its associated confidence interval based on 1000 bootstrapped samples, controlling for behavioural activity, background temperature and time held in the pen. We then averaged the readings for each individual and analysed whether this average baseline temperature, alongside fixed effects of sex, tarsus length and body condition, predicted the role (aggressor or recipient) adopted in a given encounter, using a generalized linear mixed-effects model (GLMM) in *lme4* with a binomial error function and intercepts varying across individuals. Note that whereas mass and tarsus length are highly correlated ($r = 0.81$, $t_{124} = 15.1$, $p < 0.001$), tarsus length and body condition are not ($r = -0.12$, $t_{124} = -1.3$, $p = 0.192$), which allowed us to estimate their statistical effects separately.

For the aggression thermal measurements, time was coded from -5 to 20 s, with 0 s representing the moment of attack. There were a large number of missing values due to one or both individuals briefly disappearing from view, so (following [54]) we used linear interpolation in the package *zoo* [77] to infer the most parsimonious values between two or more recorded temperatures. For example, if 36.2°C was recorded at time 1 s and 36.8°C at 4 s, we inferred the missing temperatures at 2 and 3 s to be 36.4 and 36.6°C, respectively. Note that no missing values were replaced before the first or after the last recorded temperature for a given individual in a given encounter.

We then used two separate LMMs to model the temperature profile before (-5 to 0 s) and after (1 to 20 s) the moment of attack. These models included fixed effects of time, role (aggressor or recipient), average baseline temperature (mean of 30 s measurements for that pheasant), type of aggression (threat or peck), sex, tarsus length, body condition and group, and two random effects that allowed intercepts to vary both across individual pheasants and across the contests they were involved in. We modelled nonlinear changes using polynomial terms for time and allowed this relationship to differ between aggressors and recipients by including a role \times time interaction term, but these more complex terms were omitted where they did not significantly improve the fit of the model. We also checked whether there was any difference between aggressors and recipients in the timing of the highest and lowest temperatures during an encounter, using approximate two-sample Fisher–Pitman permutation tests in the package *coin* [78]. For visualization and to check the robustness of our conclusions, we fitted a generalized additive mixed-effects model (GAMM) across the whole time sequence (-5 to 20 s) in the package *brms* [79], with smooth terms for the nonlinear effect of time on temperature.

For all mixed-effects models, we used the package *DHARMA* [80] to generate diagnostic plots of the residuals, which revealed no strong departures from the assumptions of normality and homoscedasticity. The significance of fixed effects was computed using likelihood-ratio tests, comparing the residual deviance of models including versus omitting that predictor. Continuous predictors with a zero value outside the range of measurement (i.e. morphometric and temperature variables) were standardized before analysis to aid interpretation.

The data files and associated R script are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.w0vt4b8rj> [81].

3. Results

(a) Baseline measurements: is head temperature individually repeatable?

We obtained 610 baseline head temperature measurements at 30 s intervals for 94 pheasants (1 to 18 measurements




Figure 2. Boxplots showing the distribution (median, interquartile range and outliers) of maximum head temperatures (averaged within individuals) of male and female captive juvenile pheasants engaged in different behavioural activities. (Online version in colour.)

Table 3. Estimated fixed effects in an LMM predicting maximum head temperature (°C) of captive juvenile pheasants measured every 30 s (baseline measurements), with random effects of individual pheasant ($n = 94$). Significant effects are shown in bold.

fixed effect	estimate \pm s.e.	χ^2_{c}	d.f.	p
intercept ^a	35.842 \pm 0.312			
behavioural activity		17.7	5	0.003
resting	0.862 \pm 0.285			
standing	0.606 \pm 0.247			
preening	0.578 \pm 0.292			
foraging	0.678 \pm 0.284			
walking	0.303 \pm 0.258			
background temperature^b	0.597 \pm 0.218	7.6	1	0.006
time held in pen	-0.002 \pm 0.014	<0.1	1	0.848
group		18.1	3	<0.001

^aPredicted maximum head temperature for a pheasant engaged in 'other' activities at the start of the observation session at the average background temperature.

^bStandardized before analysis (z-score method).

^cChange in deviance from a likelihood-ratio test comparing models that include or omit that predictor.

per pheasant; median = 6 measurements), excluding an additional 64 measurements that occurred within 20 s of an aggressive encounter involving the focal pheasant. The maximum temperature identified by *FLIR Tools* had an average value (mean \pm s.d.) across all measurements of 36.63 \pm 1.04°C (range 31.9–39.1°C), compared to a background temperature (white plastic drinker filled with water) of 25.82 \pm 1.46°C (range 23.2–27.5°C).

An LMM with intercepts varying across individuals showed that maximum head temperature was positively associated with background temperature and dependent on behaviour (table 3), with lower head temperatures when walking compared to resting (figure 2). Controlling for these effects, maximum head temperature showed low but highly

significant repeatability within individual pheasants (between-individual variance $\alpha^2 = 0.159$, within-individual variance $\alpha^2 = 0.873$; repeatability $R_M = 0.154$, 95% CI = 0.08–0.24; $\chi^2 = 30.6$, $p < 0.001$). Inclusion of individual-level variables in the model (which reduced repeatability to $R_{M,adj} = 0.116$, 95% CI = 0.05–0.20) showed that this was partly due to a sex difference, with males having higher maximum head temperatures on average than females ($b \pm$ s.e. = 0.472 \pm 0.158, $\chi^2 = 9.2$, $p = 0.002$; figure 2), whereas there was no relationship with morphological traits at release (tarsus length: $\chi^2 = 0.4$, $p = 0.520$; body condition: $\chi^2 = 1.0$, $p = 0.326$).

(b) Does average baseline temperature predict aggressor and recipient roles?

We observed 85 pairwise aggressive encounters (21 threats and 64 pecks, with fighting never observed) in which one or both individuals could be identified from their numbered tags. A GLMM with a binomial error structure and intercepts varying across individuals showed that the role (aggressor or recipient) adopted in a given contest was unrelated to size (tarsus length at release: odds ratio = 1.78, 95% CI = 0.72–4.63; $\chi^2 = 1.6$, $p = 0.200$), body condition (odds ratio = 1.17, 95% CI = 0.58–2.30; $\chi^2 = 0.2$, $p = 0.628$) or sex, with males ($n = 34$) and females ($n = 21$) equally likely to be the aggressor (odds ratio = 2.56, 95% CI = 0.63–13.48; $\chi^2 = 1.8$, $p = 0.186$). The average of an individual's baseline temperature measurements was very similar for aggressors (mean \pm s.e. = 36.64 \pm 0.08°C) and recipients (36.49 \pm 0.07°C) and did not reliably predict whether they would be the aggressor in a given encounter (odds ratio = 1.32, 95% CI = 0.77–2.42; $\chi^2 = 1.05$, $p = 0.306$).

(c) Aggression measurements: does head temperature change during an aggressive encounter?

There were 82 pairwise aggressive encounters, involving 55 individually identified pheasants (34 males, 21 females), for which we could measure the maximum head temperature of either the aggressor ($n = 23$), the recipient ($n = 40$) or both ($n = 19$) at some point during the encounter. These encounters gave a total of $n = 906$ temperature measurements, which we increased to $n = 1850$ using linear interpolation at 1 s intervals between two or more observed values.

The temperature profile during an aggressive encounter showed a strongly nonlinear pattern, showing a slight drop prior to the attack and then increasing sharply afterwards, before falling back down towards the baseline (figure 3). The lowest recorded temperature was very similar for aggressors (mean \pm s.e. = 35.67 \pm 0.11°C) and recipients (35.67 \pm 0.06°C) and occurred at a similar time point for both roles (permutation test: $z = 0.24$, $p = 0.817$). Likewise, the highest recorded temperature was very similar for aggressors (37.67 \pm 0.10°C) and recipients (37.45 \pm 0.09°C), and there was no clear difference in its timing ($z = 0.19$, $p = 0.857$).

Temperature dropped significantly leading up to the moment of the attack (time -5 to 0 s; table 4), but this drop did not differ between aggressors and recipients (LMM, role \times time interaction term: $\chi^2 = 0.6$, $p = 0.435$). After an attack (time 1 to 20 s), temperature increased and then decreased (significant quadratic term; table 5), but there was no clear difference between aggressors and recipients in this nonlinear pattern (LMM, joint contribution of role \times




Figure 3. Change in head temperatures during an aggressive encounter between captive juvenile pheasants in aggressor and recipient roles. Lines are conditional smooths with 95% uncertainty intervals, generated from a GAMM using the package *brms* [79]. Boxplots show the distribution (median, interquartile range and outliers) of the times (averaged within individuals within encounters) at which the minimum (bottom) and maximum (top) temperatures occurred and their observed values (right). (Online version in colour.)

Table 4. Estimated fixed effects in an LMM predicting maximum head temperature (°C) of captive juvenile pheasants leading up to an aggressive encounter (aggressive measurements, -5 to 0 s prior to the moment of attack), with random effects of individual pheasant ($n = 52$) and encounter ($n = 71$). Significant effects are shown in bold.

fixed effect	estimate \pm s.e.	χ^2 ^c	d.f.	<i>p</i>
intercept ^a	36.507 \pm 0.212			
time	-0.047 \pm 0.017	7.7	1	0.006
role (aggressor)	-0.042 \pm 0.109	0.2	1	0.691
average baseline temperature ^b	0.135 \pm 0.078	3.6	1	0.057
sex (female)	-0.344 \pm 0.203	3.2	1	0.075
tarsus length ^b	-0.116 \pm 0.127	0.9	1	0.353
body condition ^b	-0.102 \pm 0.091	1.4	1	0.245
type of encounter (threat)	-0.189 \pm 0.165	1.3	1	0.254
group		0.8	3	0.842

^aPredicted maximum head temperature at time 0 (moment of attack) for a male pheasant (recipient) of average size and with average baseline temperature, being pecked by another pheasant (aggressor).

^bStandardized before analysis (z-score method).

^cChange in deviance from a likelihood-ratio test comparing models that include or omit that predictor.

time and role \times time² interaction terms: $\chi_2^2 = 2.4$, $p = 0.299$). Overall, across the 20 s following an attack, aggressors were 0.2°C hotter than recipients and females were 0.4°C cooler than males (table 5).

To check the consistency of our results, we also modelled temperature changes across the whole time course of the encounter (time -5 to 20 s) using smooth terms rather than a polynomial function (figure 3). Our conclusions from this approach were the same: the smooth term was clearly non-linear (GAMM: estimated variance parameter = 1.42, 95% CI 0.54–2.87) and gave a substantially better fit than a linear term (Δ elpd-LOO = 29.6), but separating this smooth term

Table 5. Estimated fixed effects in an LMM predicting maximum head temperature (°C) of captive juvenile pheasants following an aggressive encounter (aggressive measurements, 1 to 20 s after the moment of attack), with random effects of individual pheasant ($n = 53$) and encounter ($n = 77$). Significant effects are shown in bold.

fixed effect	estimate \pm s.e.	χ^2 ^c	d.f.	<i>p</i>
intercept ^a	36.733 \pm 0.174			
time				
linear	0.051 \pm 0.011	22.0	1	<0.001
quadratic	-0.003 \pm 0.001	29.2	1	<0.001
role (aggressor)	0.186 \pm 0.091	4.3	1	0.037
average baseline temperature^b	0.123 \pm 0.065	4.2	1	0.040
sex (female)	-0.424 \pm 0.178	6.3	1	0.012
tarsus length ^b	-0.136 \pm 0.110	1.7	1	0.188
body condition ^b	-0.079 \pm 0.077	1.3	1	0.255
type of encounter (threat)	-0.168 \pm 0.116	1.7	1	0.194
group		0.7	3	0.872

^aPredicted maximum head temperature at time 0 (moment of attack) for a male pheasant (recipient) of average size and with average baseline temperature, being pecked by another pheasant (aggressor).

^bStandardized before analysis (z-score method).

^cChange in deviance from a likelihood-ratio test comparing models that include or omit that predictor.

between aggressors and recipients did not improve the model (Δ elpd-LOO = 1.4) [81].

As a further check on the robustness of our results, we repeated the analysis on a reduced dataset ($n = 906$) with the interpolated points removed. The effect estimates from this analysis were very similar to those from our original analysis (full results in electronic supplementary material); in particular, maximum head temperature dropped significantly leading up to the moment of attack ($b \pm$ s.e. = -0.058 ± 0.025 ,

$\chi_1^2 = 5.4, p = 0.021$; electronic supplementary material, table S1), while after the attack it rose and then fell (linear term: 0.063 ± 0.021 , $\chi_1^2 = 8.7, p = 0.003$; quadratic term: -0.003 ± 0.001 , $\chi_1^2 = 8.3, p = 0.004$; electronic supplementary material, table S2).

4. Discussion

In this study, we monitored thermographic changes in captive flocks of juvenile pheasants while they engaged in spontaneous aggressive interactions during a brief period of confinement. We found that head temperature dropped sharply in the few seconds prior to an attack, followed by an increase and then a more gradual decline back down towards baseline levels. Aggressors were on average slightly hotter than recipients, but the changes in temperature were similar for both roles. These findings are based on a novel application of IRT, which is an increasingly popular technique to study SIH in birds and other endotherms [32,49–53,55,58,59,61]. The ability to measure such physiological responses without any physical contact with the animals makes IRT potentially applicable to a wide range of systems, giving a clear advantage over other techniques that rely on implanted or wearable devices [31,46]. Previous avian IRT work has largely focused on the responses to acute non-social stressors [32,51] or long-term exposure to adverse health or welfare conditions [35,60,61], overlooking immediate short-term responses to the social interactions that establish and maintain dominance hierarchies. To the best of our knowledge, our study is the first to show detectable changes in body surface temperature during individual aggressive events occurring spontaneously in a group setting.

(a) Baseline measurements

Our baseline temperature measurements, taken every 30 s, showed low but highly significant repeatability within individual pheasants. This was partly attributable to a sex difference (females were on average $0.4\text{--}0.5^\circ\text{C}$ cooler than males), but even after accounting for this effect there were consistent individual differences in maximum head temperature, which may reflect underlying differences in metabolic rate [82] and potentially stress physiology [52,66]. An interesting avenue for future work would be to explore the extent to which these differences are linked to genotype, by comparing the temperature profiles of related and unrelated individuals. We did not find evidence that temperature differences were related to behavioural role: average baseline temperature did not predict whether an individual would be the aggressor or the recipient in a given dominance encounter. More extensive observations of these pheasants would be needed to establish whether any behavioural differences are linked to repeatable differences in physiology, as has been suggested elsewhere [67,83]. Furthermore, given that our measurements were taken from a single 13 min period for each group of pheasants, it is unclear whether the observed temperature differences would persist over a longer timescale, or are more reflective of transient differences in state on a particular day. It would be interesting to investigate this by collecting equivalent measurements across multiple days.

Despite the significant repeatability, we also found that maximum head temperature varied within individuals depending on their behavioural state, as reported previously in chickens [32]. Temperatures appeared to be lowest in more

active states such as walking and foraging compared to resting, perhaps because greater activity redirects more peripheral blood towards the muscle tissue where it is needed. This finding is consistent with human studies showing that skin surface temperature falls with increasing exercise intensity, mediated by cutaneous vasoconstriction [84].

(b) Aggression measurements

During aggressive encounters with another pheasant, our per-second measurements revealed a rapid change in maximum head temperature both before and after the moment of attack. In the 5 s leading up to an attack, we observed a drop in surface temperature that likely reflects cutaneous vasoconstriction and the redirection of peripheral blood towards the body core, causing a concurrent increase in core body temperature [31]. This physiological change is consistent with the core hyperthermic response to social defeat seen in laboratory studies of rodents [40–42] and passerines [43], commonly measured using implanted devices or invasive probes. Here, however, we detected a thermal response non-invasively during spontaneous, brief and acute dominance encounters between group mates, which were unlike the more severe and asymmetric aggression artificially induced by the resident–intruder paradigm, where a larger or highly aggressive resident individual delivers attacks on a smaller or submissive intruder placed within its home cage [44].

Having dropped prior to the attack, maximum head temperature increased to a peak, then steadily decreased towards baseline levels over the 20 s afterwards. This non-linear pattern is remarkably similar in shape (though on a shorter timescale) to that observed in two different avian IRT studies involving a handling stressor. Domestic chickens showed an initial drop of $1.3\text{--}2.2^\circ\text{C}$ in wattle and comb temperature when put in a ‘side-pinned’ hold (presumed to be a severe acute stressor), followed by a significant increase above baseline more than 10 min after release [50]. In wild blue tits (*Cyanistes caeruleus*), mean eye region surface temperature dropped by around 1°C within 10 s of being suddenly trapped inside a box, then increased to a peak around 1°C above baseline when they were captured, held in the hand and blood sampled by an experimenter, followed by a decline back towards the baseline lasting for 1–2 min after capture [54]. The reason for this temperature peak after the initial drop is unclear, but it may be linked to the mechanisms that re-establish homeostasis after SIH [31]. Intriguingly, a temperature peak was not observed in chickens that were cradled rather than side-pinned [50], nor in blue tits that were trapped but not handled or blood sampled [52]. The fact that we observed such a peak in freely interacting pheasants shows that transfer of heat from the experimenter’s hands [54] or a specific mode of handling cannot be the explanation in this case. An alternative possibility is that the post-drop peak is a response to more intense stressors [50,54], which would support the notion that dominance interactions are a highly salient, albeit short-lived, stimulus for these group-living birds. It is also possible that the temperature changes we observed were linked to rapid head movements by both birds, as the aggressor attempted to peck the recipient and the recipient attempted to avoid being pecked. More research is needed to identify the particular circumstances that induce this characteristic nonlinear pattern and to understand the physiological mechanisms driving it.

The temperature changes we recorded were rapid, with the biggest differences generally seen within 5 s either side of the moment of attack (figure 3). The fact that in our study the temperature drop began before the moment of attack suggests that there was some anticipation of the impending aggression, by both aggressors and recipients. On several occasions, we observed that the aggressor and recipient were directly facing each other and the aggressor moved towards the recipient before delivering a peck or threat, in which case both birds would have been aware of the other's presence and potentially able to prepare for an imminent attack. We also observed some bursts of aggression [85,86] in which a series of attacks spread quickly through the flock, in which case there may have been more general anticipatory responses to the occurrence of aggression nearby. After the attack, aggressors and recipients usually moved quickly away from each other, which perhaps explains why the post-stressor temperature changes were more rapid than those seen in chickens [50] and blue tits [52,54] during experimental procedures lasting a minute or more.

We predicted that the recipients of aggressive behaviour would show a stronger thermographic response than the aggressors, based on the expectation that being threatened or physically attacked would be a more 'stressful' experience than choosing to deliver that aggression. Yet there was limited evidence to support this prediction. The maximum head temperature of recipients was 0.2°C lower than that of aggressors across the 20 s following an attack, but although this difference was significant, the predominant pattern was that recipients and aggressors showed the same, strongly nonlinear change in temperature. This finding indicates that, rather than acting as a stressor for one party but not the other, directed aggressive interactions may in fact induce similar physiological responses in the aggressor and the recipient, at least in the short term. Numerous other studies support this interpretation, suggesting that the vertebrate neuroendocrine 'stress' response (via the sympatho-adrenomedullary and HPA systems) actually reflects metabolic and cardiovascular demands associated with behavioural activity, rather than the rewarding or aversive nature of the triggering stimulus [68,87]. For example, glucocorticoid levels were elevated in both the winners and losers of aggressive conflicts in rainbow trout [25], green anoles [26] and laboratory rats (*Rattus norvegicus*) [68], while winning and losing rats also showed similar peak responses in heart rate, blood pressure and core body temperature [68]. The main difference between winners and losers observed in these studies was in the speed of return to baseline (pre-conflict) values, with losers showing more prolonged elevation of physiological parameters [25,26,68]. In the pheasants, we did not find such a difference between aggressors and recipients, although this may reflect the relatively mild nature of the social defeats that were inflicted in our set-up, compared to those in experimental rodent studies. To allow a proper comparison, it would be interesting to take IRT measurements from animals involved in aggressive encounters under more controlled conditions (e.g. the resident-intruder paradigm) and follow them over a longer period of time (e.g. up to an hour afterwards).

(c) Limitations and future directions

Our study represents a first step towards characterizing the short-term physiological responses to social competition,

but there are of course some limitations. The main weakness is a lack of standardization in our measurements: unlike some IRT studies using non-social stressors [54,55], we were unable to control physical parameters such as the distance and angle of the focal bird relative to the camera, which are known to influence thermographic measurements [88]. This issue is partly mitigated by the fact that our main analysis concerned within-individual changes in temperature over a brief period (25 s) during which both birds (aggressor and recipient) would have been in a similar part of the pen. There were also many varying aspects of the social context (e.g. aggression between other birds close by) that we did not attempt to control, which will have added statistical noise to our measurements compared to more standardized protocols (e.g. staged encounters in the resident-intruder paradigm). Given such limitations, it is promising that in this noisy environment we could still detect a clear thermographic response to aggressive encounters comparable to that seen for non-social stressors [50,54], but taking similar measurements in a more controlled setting would help to validate our findings. A more standardized protocol would also allow measurements to be collected over a longer period after the aggressive attack, to establish whether recipients have a slower return to their baseline surface temperature than aggressors, as observed for other physiological parameters [25,26,68].

At the same time, the lack of experimental control is an important strength of our study, lending our results greater ecological validity. The acts of aggression we observed arose spontaneously among familiar, same-aged birds that had been reared together since hatching, as is standard practice in the gamebird industry [89]. Captive pheasants are aggressive towards one another within the first few weeks of life [65] and dominance relationships can continue to change as males move into adulthood [90], so the behaviour we recorded likely represents the early emergence of a dominance hierarchy that can ultimately shape their sexual success [64]. By contrast, the resident-intruder paradigm commonly used in rodent studies of aggression is a more controlled but contrived setting, where the intruder is forcibly exposed to an aggressive resident, with the odds stacked heavily against the former [44]. Another strength is that the birds in our study engaged in aggressive behaviour while surrounded by their flock mates, potentially allowing bystander effects [12] to operate. Detailed analysis of dominance hierarchies in other galliforms suggests that they are acutely sensitive to interactions between other individuals in the flock, and that this affects hierarchy formation [85]. For these reasons, although the resident-intruder paradigm has clearly provided valuable insights into the physiological consequences of social defeat, we suggest that it sheds less light on the establishment and maintenance of dominance in groups of freely interacting individuals.

There are a number of useful ways that future studies could build on our findings. In particular, it would be valuable to combine IRT with a more dynamic view of dominance [91] than the snapshot presented here. One obvious extension would be to collect thermographic measurements across a longer period spanning several weeks, as the dominance hierarchy develops. As well as enabling a clearer assessment of the individual repeatability of thermal profiles, this would reveal any changes as individuals ascend or descend in dominance rank. In our preliminary observations here we found that aggression was never met with retaliation and thus the aggressors were always victors, in line with evidence from chickens

that ‘pair-flips’ are relatively rare [3,85]. Over a longer time-scale, however, we would expect to see some dominance relationships being reversed. It would be particularly interesting to examine the thermographic changes during such dominance reversals, as arguably these should induce stronger physiological responses than encounters where the direction of dominance is maintained. Similarly, we might expect to see bigger thermographic changes when conflicts escalate to physical fighting, which was never observed in our study but is known to happen over longer periods [63]. Such effects might extend to individuals not directly involved in the dominance encounter, as has been reported for greylag geese using implanted heart rate monitors [45]. By taking thermographic measurements from all flock members when such escalated interactions occur, it might be possible to detect physiological correlates of bystander effects. IRT could also be used to compare physiological responses between different types of behavioural interactions that are used strategically in different contexts [92], as well as to compare strategic with non-strategic interactions. Another worthwhile follow-up study would be to conduct a more fine-grained analysis of the behavioural movements that occur during aggressive interactions. For example, it would be interesting to quantify the head movements of aggressors when attempting to peck recipients, and those of the recipients as they attempt to avoid being pecked, to examine whether the pattern of movement predicts individual changes in head temperature. Finally, as a complementary approach to all of these suggestions, it would be valuable to take thermographic measurements from interacting animals fitted with implanted devices, so that the data can be compared and validated against other, more established indicators of physiological arousal (e.g. heart rate [46] and core body temperature [31]).

A century has passed since Schjelderup-Ebbe’s landmark paper on pecking orders in chickens [3]. Alongside his pioneering behavioural observations he made several intriguing comments about the birds’ emotional state in the face of social challenges, referring to their ‘anger’, ‘fear’ and

‘courage’ (translation of [3] by M. Schleidt & W. M. Schleidt), despite not having any data on physiological variables, let alone cognitive or emotional appraisal. A proliferation of sampling techniques in recent decades has provided many windows on the physiological response to dominance conflicts and other stressors [27], encouraging a richer view of dominance than one based purely on overt aggressive behaviour [93]. Our data on groups of freely interacting individuals suggest that IRT can offer additional valuable insights into the mechanisms driving the formation of animal dominance hierarchies.

Ethics. All work was approved by the University of Exeter Psychology Research Ethics Committee and formed part of a larger research programme conducted under UK Home Office licence PPL 30/3204 (issued to J.R.M.). Husbandry procedures adhered to the code of practice of the UK Department for Environment, Food & Rural Affairs [94], with rearing densities lower than recommended limits. The pheasants were held in their indoor pen for no longer than 30 min and no injuries were observed during filming.

Data accessibility. The datasets supporting this article are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.w0vt4b8rj> [81].

The data are provided in electronic supplementary material [95].

Authors’ contributions. T.W.F. and J.R.M. conceived the study. All authors helped to design the methodology. S.K. collected the behavioural and thermographic data, while M.A.W. and J.R.M. collected the morphometric data. T.W.F. and S.K. analysed the data. S.K. and T.W.F. wrote the manuscript with input from all other authors.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Competing interests. We declare we have no competing interests.

Funding. This work was funded by the European Research Council (grant no. 616474 to J.R.M.) and the Association for the Study of Animal Behaviour (ASAB Research Grant to P.E.R.).

Acknowledgments. We thank Eli Strauss, Liz Hobson, Dai Shizuka and James Curley for organizing this special issue, two anonymous referees for valuable feedback, and Doretta Caramaschi and members of the CRAB group at Exeter for helpful discussion. Christine Beardsworth, Kandace Griffin, Pip Laker, Anna Morris and Jayden van Horik assisted with the collection of morphometric data and general care of the birds.

References

- Drews C. 1993 The concept and definition of dominance in animal behavior. *Behaviour* **125**, 283–313. (doi:10.1163/156853993X00290)
- Clutton-Brock TH, Huchard E. 2013 Social competition and selection in males and females. *Phil. Trans. R. Soc. B* **368**, 20130074. (doi:10.1098/rstb.2013.0074)
- Schjelderup-Ebbe T. 1922 Beitrage zur Sozialpsychologie des Haushuns. *Z. Psychol.* **88**, 225–252.
- Hobson EA. 2022 Quantifying the dynamics of nearly 100 years of dominance hierarchy research. *Phil. Trans. R. Soc. B* **377**, 20200433. (doi:10.1098/rstb.2020.0433)
- Tibbetts EA, Pardo-Sanchez J, Weise C. 2022 The establishment and maintenance of dominance hierarchies. *Phil. Trans. R. Soc. B* **377**, 20200450. (doi:10.1098/rstb.2020.0450)
- Favati A, Leimar O, Løvlie H. 2014 Personality predicts social dominance in male domestic fowl. *PLoS ONE* **9**, e103535. (doi:10.1371/journal.pone.0103535)
- Pottinger TG, Carrick TR. 2001 Stress responsiveness affects dominant–subordinate relationships in rainbow trout. *Horm. Behav.* **40**, 419–427. (doi:10.1006/hbeh.2001.1707)
- Chase ID, Tovey C, Spangler-Martin D, Manfredonia M. 2002 Individual differences versus social dynamics in the formation of dominance hierarchies. *Proc. Natl Acad. Sci. USA* **99**, 5744–5749. (doi:10.1073/pnas.082104199)
- Chase ID, Bartolomeo C, Dugatkin LA. 1994 Aggressive interactions and inter-contest interval: how long do winners keep winning? *Anim. Behav.* **48**, 393–400. (doi:10.1006/anbe.1994.1253)
- Hsu Y, Earley RL, Wolf LL. 2006 Modulation of aggressive behaviour by fighting experience: mechanisms and contest outcomes. *Biol. Rev.* **81**, 33–74. (doi:10.1017/S146479310500686X)
- Rutte C, Taborsky M, Brinkhof MWF. 2006 What sets the odds of winning and losing? *Trends Ecol. Evol.* **21**, 16–21. (doi:10.1016/j.tree.2005.10.014)
- Dugatkin LA. 2001 Bystander effects and the structure of dominance hierarchies. *Behav. Ecol.* **12**, 348–352. (doi:10.1093/beheco/12.3.348)
- Qu C, Ligneul R, Van der Henst JB, Dreher JC. 2017 An integrative interdisciplinary perspective on social dominance hierarchies. *Trends Cogn. Sci.* **21**, 893–908. (doi:10.1016/j.tics.2017.08.004)
- Creel S. 2001 Social dominance and stress hormones. *Trends Ecol. Evol.* **16**, 491–497. (doi:10.1016/S0169-5347(01)02227-3)
- Sapolsky RM. 2005 The influence of social hierarchy on primate health. *Science* **308**, 648–652. (doi:10.1126/science.1106477)

16. Goymann W, Wingfield JC. 2004 Allostatic load, social status and stress hormones: the costs of social status matter. *Anim. Behav.* **67**, 591–602. (doi:10.1016/j.anbehav.2003.08.007)
17. Rubenstein DR, Shen SF. 2009 Reproductive conflict and the costs of social status in cooperatively breeding vertebrates. *Am. Nat.* **173**, 650–661. (doi:10.1086/597606)
18. Creel S, Dantzer B, Goymann W, Rubenstein DR. 2013 The ecology of stress: effects of the social environment. *Funct. Ecol.* **27**, 66–80. (doi:10.1111/j.1365-2435.2012.02029.x)
19. Williamson CM, Lee W, Romeo RD, Curley JP. 2017 Social context-dependent relationships between mouse dominance rank and plasma hormone levels. *Physiol. Behav.* **171**, 110–119. (doi:10.1016/j.physbeh.2016.12.038)
20. McCowan B, Vandeleeest J, Balasubramaniam K, Hsieh F, Nathman A, Beisner B. 2022 Measuring dominance certainty and assessing its impact on individual and societal health in a nonhuman primate model: a network approach. *Phil. Trans. R. Soc. B* **377**, 20200438. (doi:10.1098/rstb.2020.0438)
21. Milewski TM, Lee W, Champagne FA, Curley JP. 2022 Behavioural and physiological plasticity in social hierarchies. *Phil. Trans. R. Soc. B* **377**, 20200443. (doi:10.1098/rstb.2020.0443)
22. Simons ND, Michopoulos V, Wilson M, Barreiro LB, Tung J. 2022 Agonism and grooming behaviour explain social status effects on physiology and gene regulation in rhesus macaques. *Phil. Trans. R. Soc. B* **377**, 20210132. (doi:10.1098/rstb.2021.0132)
23. Summers CH, Watt MJ, Ling TL, Forster GL, Carpenter RE, Korzan WJ, Lukkes JL, Øverli Ø. 2005 Glucocorticoid interaction with aggression in non-mammalian vertebrates: reciprocal action. *Eur. J. Pharmacol.* **526**, 21–35. (doi:10.1016/j.ejphar.2005.09.059)
24. Summers CH, Winberg S. 2006 Interactions between the neural regulation of stress and aggression. *J. Exp. Biol.* **209**, 4581–4589. (doi:10.1242/jeb.02565)
25. Øverli Ø, Harris CA, Winberg S. 1999 Short-term effects of fights for social dominance and the establishment of dominant–subordinate relationships on brain monoamines and cortisol in rainbow trout. *Brain Behav. Evol.* **54**, 263–275. (doi:10.1159/00006627)
26. Summers CH, Summers TR, Moore MC, Korzan WJ, Woodley SK, Ronan PJ, Höglund E, Watt MJ, Greenberg N. 2003 Temporal patterns of limbic monoamine and plasma corticosterone response during social stress. *Neuroscience* **116**, 554–563. (doi:10.1016/S0306-4522(02)00708-X)
27. Gormally BMG, Romero ML. 2020 What are you actually measuring? A review of techniques that integrate the stress response on distinct time-scales. *Funct. Ecol.* **34**, 2030–2044. (doi:10.1111/1365-2435.13648)
28. Romero LM, Reed JM. 2005 Collecting baseline corticosterone samples in the field: is under 3 min good enough? *Comp. Biochem. Physiol. A* **140**, 73–79. (doi:10.1016/j.cbpb.2004.11.004)
29. Harper JM, Austad SN. 2000 Fecal glucocorticoids: a noninvasive method of measuring adrenal activity in wild and captive rodents. *Phys. Biochem. Zool.* **73**, 12–22. (doi:10.1086/316721)
30. Gesquiere LR, Learn NH, Simao CM, Onyango PO, Alberts SC, Altmann J. 2011 Life at the top: rank and stress in wild male baboons. *Science* **333**, 357–360. (doi:10.1126/science.1207120)
31. Oka T, Oka K, Hori T. 2001 Mechanisms and mediators of psychological stress-induced rise in core temperature. *Psychosom. Med.* **63**, 476–486. (doi:10.1097/00006842-200105000-00018)
32. Edgar JL, Nicol CJ, Pugh CA, Paul ES. 2013 Surface temperature changes in response to handling in domestic chickens. *Physiol. Behav.* **119**, 195–200. (doi:10.1016/j.physbeh.2013.06.020)
33. Marazziti D, Di Muro A, Castrogiovanni P. 1992 Psychological stress and body temperature changes in humans. *Physiol. Behav.* **52**, 393–395. (doi:10.1016/0031-9384(92)90290-1)
34. Briese E. 1995 Emotional hyperthermia and performance in humans. *Physiol. Behav.* **58**, 615–618. (doi:10.1016/0031-9384(95)00091-V)
35. Iyasere OS, Edwards SA, Bateson M, Mitchell M, Guy JH. 2017 Validation of an intramuscularly-implanted microchip and a surface infrared thermometer to estimate core body temperature in broiler chickens exposed to heat stress. *Comput. Electron. Agric.* **133**, 1–8. (doi:10.1016/j.compag.2016.12.010)
36. Kramer K, van de Weerd H, Mulder A, van Heijningen C, Baumans V, Remie R, Voss HP, van Zutphen BFM. 2004 Effect of conditioning on the increase of heart rate and body temperature provoked by handling in the mouse. *Altern. Lab. Anim.* **32**, 177–181. (doi:10.1177/026119290403201s29)
37. Meijer MK, Spruijt BM, van Zutphen LFM, Baumans V. 2006 Effect of restraint and injection methods on heart rate and body temperature in mice. *Lab. Anim.* **40**, 382–391. (doi:10.1258/002367706778476370)
38. Caramaschi D, de Boer SF, Koolhaas JM. 2007 Differential role of the 5-HT1A receptor in aggressive and non-aggressive mice: an across-strain comparison. *Physiol. Behav.* **90**, 590–601. (doi:10.1016/j.physbeh.2006.11.010)
39. Caramaschi D. 2009 *The physiology of aggression: towards understanding violence*. PhD thesis, University of Groningen, The Netherlands.
40. Sgoifo A, Pozzato C, Meerlo P, Costoli T, Manghi M, Stilli D, Olivetti G, Musso E. 2002 Intermittent exposure to social defeat and open-field test in rats: acute and long-term effects on ECG, body temperature and physical activity. *Stress* **5**, 23–35. (doi:10.1080/102538902900012387)
41. Keeney AJ, Hogg S, Marsden CA. 2001 Alterations in core body temperature, locomotor activity, and corticosterone following acute and repeated social defeat of male NMRI mice. *Physiol. Behav.* **74**, 177–184. (doi:10.1016/S0031-9384(01)00541-8)
42. Bartolomucci A, Palanza P, Costoli T, Savani E, Laviola G, Parmigiani S, Sgoifo A. 2003 Chronic psychosocial stress persistently alters autonomic function and physical activity in mice. *Physiol. Behav.* **80**, 57–67. (doi:10.1016/S0031-9384(03)00209-9)
43. Carere C, Welink D, Drent PJ, Koolhaas JM, Groothuis TGG. 2001 Effect of social defeat in a territorial bird (*Parus major*) selected for different coping styles. *Physiol. Behav.* **73**, 427–433. (doi:10.1016/S0031-9384(01)00492-9)
44. Martinez M, Calvo-Torrent A, Pico-Alfonso MA. 1998 Social defeat and subordination as models of social stress in laboratory rodents: a review. *Aggress. Behav.* **24**, 241–256. (doi:10.1002/(SICI)1098-2337(1998)24:4<241::AID-AB1>3.0.CO;2-M)
45. Wascher CAF, Scheiber IBR, Kotrschal K. 2008 Heart rate modulation in bystanding geese watching social and non-social events. *Proc. R. Soc. B* **275**, 1653–1659. (doi:10.1098/rspb.2008.0146)
46. Wascher CAF. 2021 Heart rate as a measure of emotional arousal in evolutionary biology. *Phil. Trans. R. Soc. B* **376**, 20200479. (doi:10.1098/rstb.2020.0479)
47. Busnardo C, Tavares RF, Resstel LBM, Elias LLK, Correa FMA. 2010 Paraventricular nucleus modulates autonomic and neuroendocrine responses to acute restraint stress in rats. *Auton. Neurosci.* **158**, 51–57. (doi:10.1016/j.autneu.2010.06.003)
48. Vianna DML, Carrive P. 2005 Changes in cutaneous and body temperature during and after conditioned fear to context in the rat. *Eur. J. Neurosci.* **21**, 2505–2512. (doi:10.1111/j.1460-9568.2005.04073)
49. Speakman JR, Ward S. 1998 Infrared thermography: principles and applications. *Zoology* **101**, 224–232.
50. Herborn KA, Graves JL, Jerem P, Evans NP, Nager R, McCafferty DJ, McKeegan DEF. 2015 Skin temperature reveals the intensity of acute stress. *Physiol. Behav.* **152**, 225–230. (doi:10.1016/j.physbeh.2015.09.032)
51. Herborn KA, Jerem P, Nager R, McKeegan DEF, McCafferty DJ. 2018 Surface temperature elevated by chronic and intermittent stress. *Physiol. Behav.* **191**, 47–55. (doi:10.1016/j.physbeh.2018.04.004)
52. Jerem P, Herborn K, McCafferty D, McKeegan D, Nager R. 2015 Thermal imaging to study stress non-invasively in unrestrained birds. *J. Vis. Exp.* **105**, e53184. (doi:10.3791/53184)
53. Jerem P, Jenni-Eiermann S, Herborn K, McKeegan D, McCafferty DJ, Nager RG. 2018 Eye region surface temperature reflects both energy reserves and circulating glucocorticoids in a wild bird. *Sci. Rep.* **8**, 1907. (doi:10.1038/s41598-018-20240-4)
54. Jerem P, Jenni-Eiermann S, McKeegan D, McCafferty DJ, Nager RG. 2019 Eye region surface temperature dynamics during acute stress relate to baseline glucocorticoids independently of environmental conditions. *Physiol. Behav.* **210**, 112627. (doi:10.1016/j.physbeh.2019.112627)

55. Whiteside MA, Madden JR. Submitted Using thermography to detect arousal induced by an ephemeral and low-level frustration event.
56. Veening JG, Bouwknecht JA, Joosten HJ, Dederen PJ, Zethof TJ, Groenink L, van der Gugten J, Olivier B. 2004 Stress-induced hyperthermia in the mouse: *c-fos* expression, corticosterone and temperature changes. *Prog. Neuropsychopharmacol. Biol. Psychiatry* **28**, 699–707. (doi:10.1016/j.pnpb.2004.05.007)
57. McCafferty DJ, Gallon S, Nord A. 2015 Challenges of measuring body temperatures of free-ranging birds and mammals. *Anim. Biotelemetry* **3**, 33. (doi:10.1186/s40317-015-0075-2)
58. Kuraoka K, Nakamura K. 2011 The use of nasal skin temperature measurements in studying emotion in macaque monkeys. *Physiol. Behav.* **102**, 347–355. (doi:10.1016/j.physbeh.2010.11.029)
59. Dezechache G, Zuberbühler K, Davila-Ross M, Dahl CD. 2017 Skin temperature changes in wild chimpanzees upon hearing vocalizations of conspecifics. *R. Soc. Open Sci.* **4**, 160816. (doi:10.1098/rsos.160816)
60. Wilcox CS, Patterson J, Cheng HW. 2009 Use of thermography to screen for subclinical bumblefoot in poultry. *Poult. Sci.* **88**, 1176–1180. (doi:10.3382/ps.2008-00446)
61. Abudabos AM, Samara EM, Hussein EOS, Al-Ghadi MQ, Al-Atiyat RM. 2013 Impacts of stocking density on the performance and welfare of broiler chickens. *Ital. J. Anim. Sci.* **12**, e11. (doi:10.4081/ijas.2013.e11)
62. Porter WR, Witmer LM. 2016 Avian cephalic vascular anatomy, sites of thermal exchange, and the rete ophthalmicum. *Anat. Rec.* **299**, 1461–1486. (doi:10.1002/ar.23375)
63. Hill DA, Robertson PA. 1988 *The pheasant: ecology, management and conservation*. Oxford, UK: Blackwell Scientific Books.
64. Mateos C, Carranza J. 1999 Effects of male dominance and courtship display on female choice in the ring-necked pheasant. *Behav. Ecol. Sociobiol.* **45**, 235–244. (doi:10.1007/s002650050558)
65. Whiteside MA, van Horik JO, Langley EJG, Beardsworth CE, Laker PR, Madden JR. 2017 Differences in social preference between the sexes during ontogeny drive segregation in a precocial species. *Behav. Ecol. Sociobiol.* **71**, 103. (doi:10.1007/s00265-017-2332-2)
66. Taff CC, Schoenle LA, Vitousek MN. 2018 The repeatability of glucocorticoids: a review and meta-analysis. *Gen. Comp. Endocrinol.* **260**, 136–145. (doi:10.1016/j.ygcen.2018.01.011)
67. Carere C, Caramaschi D, Fawcett TW. 2010 Covariation between personalities and individual differences in coping with stress: converging evidence and hypotheses. *Curr. Zool.* **56**, 728–740. (doi:10.1093/czoolo/56.6.728)
68. Koolhaas JM *et al.* 2011 Stress revisited: a critical evaluation of the stress concept. *Neurosci. Biobehav. Rev.* **35**, 1291–1301. (doi:10.1016/j.neubiorev.2011.02.003)
69. Haussmann MF, Longenecker AS, Marchetto NM, Juliano SA, Bowden RM. 2012 Embryonic exposure to corticosterone modifies the juvenile stress response, oxidative stress and telomere length. *Proc. R. Soc. B* **279**, 1447–1456. (doi:10.1098/rspb.2011.1913)
70. Goerlich VC, Nätt D, Elfving M, Macdonald B, Jensen P. 2012 Transgenerational effects of early experience on behavioral, hormonal and gene expression responses to acute stress in the precocial chicken. *Horm. Behav.* **61**, 711–718. (doi:10.1016/j.yhbeh.2012.03.006)
71. Casey-Trott TM, Widowski TM. 2016 Behavioral differences of laying hens with fractured keel bones within furnished cages. *Front. Vet. Sci.* **3**, 42. (doi:10.3389/fvets.2016.00042)
72. Whiteside MA. 2015 How early rearing conditions influence behaviour and survival of pheasants released into the wild. PhD thesis, University of Exeter, UK.
73. Langley EJG, van Horik JO, Whiteside MA, Beardsworth CE, Madden JR. 2018 The relationship between social rank and spatial learning in pheasants, *Phasianus colchicus*: cause or consequence? *PeerJ* **6**, e5738. (doi:10.7717/peerj.5738)
74. R Core Team. 2018 *R: a language and environment for statistical computing, version 3.6.3*. Vienna, Austria: R Foundation for Statistical Computing, <https://www.R-project.org/>.
75. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models using *lme4*. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)
76. Stoffel MA, Nakagawa S, Schielzeth H. 2017 *rptR*: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods Ecol. Evol.* **8**, 1639–1644. (doi:10.1111/2041-210X.12797)
77. Zeileis A, Grothendieck G. 2005 *zoo: S3 infrastructure for regular and irregular time series*. *J. Stat. Softw.* **14**, 1–27. (doi:10.18637/jss.v014.i06)
78. Hothorn T, Hornik K, van de Wiel MA, Zeileis A. 2008 Implementing a class of permutation tests: the *coin* package. *J. Stat. Softw.* **28**, 1–23. (doi:10.18637/jss.v028.i08)
79. Bürkner P. 2017 *brms*: an R package for Bayesian multilevel models using Stan. *J. Stat. Softw.* **80**, 1–28. (doi:10.18637/jss.v080.i01)
80. Hartig F. 2021 *DHARMA*: residual diagnostics for hierarchical (multi-level / mixed) regression models, version 0.4.1. <http://florianhartig.github.io/DHARMA/>.
81. Knoch S, Whiteside MS, Madden JR, Rose PE, Fawcett TW. 2022 Supplementary material: R script and data files. Dryad Digital Repository. (doi:10.5061/dryad.w0vt4b8rj)
82. Nespolo RF, Franco M. 2007 Whole-animal metabolic rate is a repeatable trait: a meta-analysis. *J. Exp. Biol.* **210**, 2000–2005. (doi:10.1242/jeb.02780)
83. Biro PA, Stamps JA. 2010 Do consistent individual differences in metabolic rate promote consistent individual differences in behavior? *Trends Ecol. Evol.* **25**, 653–659. (doi:10.1016/j.tree.2010.08.003)
84. Torii M, Yamasaki M, Sasaki T, Nakayama H. 1992 Fall in skin temperature of exercising man. *Br. J. Sports Med.* **26**, 29–32. (doi:10.1136/bjism.26.1.29)
85. Lindquist WB, Chase ID. 2009 Data-based analysis of winner–loser models of hierarchy formation in animals. *Bull. Math. Biol.* **71**, 556–584. (doi:10.1007/s11538-008-9371-9)
86. Lee W, Fu J, Bouwman N, Farago P, Curley JP. 2019 Temporal microstructure of dyadic social behavior during relationship formation in mice. *PLoS ONE* **14**, e0220596. (doi:10.1371/journal.pone.0220596)
87. Buwalda B, Scholte J, de Boer SF, Coppens CM, Koolhaas JM. 2012 The acute glucocorticoid stress response does not differentiate between rewarding and aversive social stimuli in rats. *Horm. Behav.* **61**, 218–226. (doi:10.1016/j.yhbeh.2011.12.012)
88. Harrap MJM, Hempel de Ibarra N, Whitney HM, Rands SA. 2018 Reporting of thermography parameters in biology: a systematic review of thermal imaging literature. *R. Soc. Open Sci.* **5**, 181281. (doi:10.1098/rsos.181281)
89. Swarbrick O. 1985 Pheasant rearing: associated husbandry and disease problems. *Vet. Rec.* **116**, 610–617. (doi:10.1136/vr.116.23.610)
90. Mateos C, Illanes MC. 1997 Dominancia social en machos de faisán común: posible origen metodológico de las discrepancias entre estudios. *Etología* **5**, 39–58.
91. Strauss ED, Shizuka D. 2022 The dynamics of dominance: open questions, challenges and solutions. *Phil. Trans. R. Soc. B* **377**, 20200445. (doi:10.1098/rstb.2020.0445)
92. Dehnen T, Papageorgiou D, Nyaguthii B, Cheronov W, Penndorf J, Boogert NJ, Farine DR. 2022 Costs dictate strategic investment in dominance interactions. *Phil. Trans. R. Soc. B* **377**, 20200447. (doi:10.1098/rstb.2020.0447)
93. Lewis RJ. 2022 Aggression, rank and power: why hens (and other animals) do not always peck according to their strength. *Phil. Trans. R. Soc. B* **377**, 20200434. (doi:10.1098/rstb.2020.0434)
94. United Kingdom Department for Environment, Food & Rural Affairs (DEFRA). 2009 *Code of practice for the welfare of gamebirds reared for sporting purposes*. Norwich, UK: DEFRA.
95. Knoch S, Whiteside MA, Madden JR, Rose PE, Fawcett TW. 2021 Hotheaded peckers: thermographic changes during aggression among juvenile pheasants (*Phasianus colchicus*). FigShare